



**PESTICIDES IN MISSISSIPPI AIR AND RAIN: A COMPARISON BETWEEN 1995 AND 2007**

**MICHAEL S. MAJEWSKI, RICHARD H. COUPE, WILLIAM T. FOREMAN, and PAUL D. CAPEL**

*Environ Toxicol Chem.*, **Accepted Article** • DOI: 10.1002/etc.2550

**Accepted Article**

"Accepted Articles" are peer-reviewed, accepted manuscripts that have not been edited, formatted, or in any way altered by the authors since acceptance. They are citable by the Digital Object Identifier (DOI). After the manuscript is edited and formatted, it will be removed from the "Accepted Articles" Web site and published as an Early View article. Note that editing may introduce changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. SETAC cannot be held responsible for errors or consequences arising from the use of information contained in these manuscripts.

**PESTICIDES IN MISSISSIPPI AIR AND RAIN: A COMPARISON BETWEEN 1995 AND 2007**

**MICHAEL S. MAJEWSKI,\*† RICHARD H. COUPE,‡ WILLIAM T. FOREMAN,§ and PAUL D. CAPELL**

† U.S. Geological Survey, Sacramento, California, USA

‡ U.S. Geological Survey, Pearl, Mississippi, USA

§ U.S. Geological Survey, Denver, Colorado, USA

|| U.S. Geological Survey, Minneapolis, Minnesota, USA

Running title: Pesticides in Mississippi air and rain

\*Address correspondence to [majewski@usgs.gov](mailto:majewski@usgs.gov).

**Additional Supporting Information may be found in the online version of this article.**

© 2014 SETAC

Submitted 21 August 2013; Returned for Revisions 3 February 2014; Accepted 6 February 2014

**Abstract:** A variety of current-use pesticides were determined in weekly-composite air and rain samples collected during the 1995 and 2007 growing seasons in the Mississippi Delta agricultural region. Similar sampling and analytical methods allowed for direct comparison of results. Decreased overall pesticide use in 2007 relative to 1995 generally resulted in decreased detection frequencies in air and rain, but observed concentration ranges were similar between years even though the 1995 sampling site was 500 m from active fields while the 2007 sampling site was within 3 m of a field. Mean concentration of detections were sometimes greater in 2007 than in 1995 but the median values were often lower. Seven compounds in 1995 and five in 2007 were detected in  $\geq 50\%$  of both air and rain samples. Atrazine, metolachlor, and propanil were detected in  $\geq 50\%$  of the air and rain samples in both years. Glyphosate and its degradation product, AMPA, were detected in  $\geq 75\%$  of air and rain samples in 2007, but were not measured in 1995. The 1995 seasonal wet depositional flux was dominated by methyl parathion (88%) and was  $>4.5$  times the 2007 flux. Total herbicide flux in 2007 was slightly greater than in 1995, and was dominated by glyphosate. Malathion, methyl parathion, and degradation products made up most of the 2007 non-herbicide flux.

**Keywords:** Pesticides, Air, Rain, Deposition flux

## INTRODUCTION

The atmosphere is an important component of the hydrologic cycle that facilitates the widespread movement of pesticides throughout the environment. Pesticides can become airborne through volatilization and wind erosion of particles both during and after their application [1–3]. Volatilization from treated areas is a continuous process and can be a major dissipative route for many pesticides [4–6]. Airborne pesticide can be carried by wind and deposited in unintended areas by dry (gas and particle) and wet (rain and snow) deposition. Rainfall efficiently cleans the atmosphere of suspended particulate matter to which pesticides might be sorbed, and gas-phase pesticides can partition directly into a falling raindrop [7]. Deposited residues can revolatilize, reenter the atmosphere, and be transported and redeposited downwind repeatedly until they are transformed or accumulate, usually in areas with cooler climates [8–10]. Atmospheric deposition of pesticides is most likely to affect the water quality in rivers and streams during runoff events when precipitation and direct surface runoff are oftentimes the major sources of streamflow.

The timing, amount applied, and the coverage area of each pesticide, as well as the proximity of pesticide applications to the sampling site, all play major roles in whether a pesticide is present in air or rain samples and at what concentration. The results of many studies on pesticide occurrence in the environment show that generally those pesticides detected most frequently and at the highest concentrations are also those that are the most heavily used [11]. Locally high pesticide concentrations in rain and air are very seasonal, are often correlated to local use, and usually occur during the spring and summer months [12, 13]. The physical and chemical properties of each pesticide also play an important role in determining if a pesticide becomes airborne, whether it then exists primarily in the gaseous or particle phase, and how efficiently rainfall removes it from the atmosphere [14, 15].

The occurrence of organochlorine compounds, such as PCBs, DDT, and others, have been documented in the atmosphere since the 1960s [16–18]. The occurrence of other pesticide classes, like the triazine, chloroacetanilide, chlorophenoxy acid, and organophosphorus pesticides, has been documented in the United States and Canada as well [14, 15, 19, 20]. In one of the most spatially comprehensive studies of the occurrence of

selected pesticides in rain in the United States, Goolsby et al. [21] found that several triazine and chloroacetanilide herbicides were present in rainfall throughout the midwestern and northeastern United States during late spring and summer in 1990 and 1991. The estimated deposition flux for atrazine and alachlor ranged from more than 240  $\mu\text{g}/\text{m}^2/\text{yr}$  in some areas of the Midwest to less than 10  $\mu\text{g}/\text{m}^2/\text{yr}$  for the New England states. Interestingly, estimates of depositional mass for alachlor, atrazine, and metolachlor from rainfall in the Midwest represent less than 1 percent of the amounts applied for each of these compounds [21–23].

The present study was conducted in northwestern Mississippi near the Mississippi River within the alluvial plain area referred to as the Delta (Figure 1). The Delta is an agriculturally intensive area where over 80% of the harvested acreage in Mississippi takes place (E Young, US Department of Agriculture National Agricultural Statistics Service, Little Rock, AR, USA, personal communication), and a broad variety of pesticides are used. From 1995 to 2007, agriculture underwent many dramatic changes that potentially affected the occurrence of pesticides in the environment. The amount of land used for agricultural production in the Delta did not change much between the 2 years. The same crops types were grown, however the proportions of some crops changed. For example, corn acreage increased from 1.21 million ha in 1995 to 3.76 million ha in 2007, whereas cotton decreased from 5.91 to 2.67 million ha during the same time period (<http://quickstats.nass.usda.gov/results/658C3944-93EB-344A-AF98-11D723DEB2CE>). Also over this time period, new developments in pest management changed how some pests, both weeds and insects, were controlled. Voluntary and regulatory decisions resulted in some pesticides being removed from the U.S. market, e.g., cyanazine was cancelled in Dec. 1999 ([http://www.epa.gov/pesticides/reregistration/status\\_page\\_c.htm](http://www.epa.gov/pesticides/reregistration/status_page_c.htm)), molinate was voluntarily phased out by 2009 (<http://www.epa.gov/fedrgstr/EPA-PEST/2008/July/Day-30/p17475.htm>), and new pesticides, such as the insecticide fipronil, were introduced or more broadly applied.

The largest single change has been the development and introduction of genetically-modified (GM) crops, particularly corn, cotton, and soybeans. These crops were introduced in the mid-1990s with genetic modifications that made them resistant to specific herbicides, or that imparted insecticidal properties within the plant itself, or both. Two of the more common modifications are (a) the resistance to glyphosate, which allows this herbicide to be applied directly to the crop for weed control throughout the growing season without harming the crop itself, and (b) having the gene codes incorporated into the plant DNA that cause the expression of *Bacillus thuringiensis* ( $B_t$ )

protein toxins for controlling certain types of insects [24]. Widespread use of these GM crops has caused major changes in the amount and type of pesticides applied to the landscape. Many herbicides previously vital to weed control in the Delta have been replaced by glyphosate [25]. In addition, the use of insecticides, including methyl parathion, in the Delta region has declined because of the increased use of GM crops incorporating B<sub>t</sub> insect control, particularly cotton and corn. These changes in pest control, and the resultant potential effects on the environment, may be similar in other areas of the country where GM crops are grown.

This present study documents the occurrence, concentration, and depositional amounts of pesticides in air and rain during the 1995 and 2007 growing seasons in the Mississippi Delta. These data are compared to changes in cropping patterns, chemical use, sampling site location and proximity to fields, to help understand pesticide movement in the atmosphere.

## **METHODS**

Weekly composite air and rainfall samples were collected during the growing season (March through September) in 1995 and 2007. The sampling sites were located in the Delta area of the lower Mississippi River watershed in west central Mississippi (Figure 1). The 1995 sampling site was located at the center of a catfish farm pond complex near the town of Rolling Fork, Mississippi and was approximately 500 m from the nearest agricultural fields [26]. This site was selected to minimize the influence of pesticide applications on nearby cotton, corn, alfalfa, and soybean fields. The 2007 sampling site was located near Pace, Mississippi approximately 100 km north of the 1995 site and within approximately 3 m of a soybean field in an area surrounded by both soybean and rice fields [27, 28]. Detailed descriptions of the methods used to collect the weekly composite air (operationally-defined gas and particle phases) and wet-only precipitation samples are given in Foreman et al. [29] and Majewski et al. [30], respectively.

Air and rain sampling for glyphosate and its degradation product aminomethyl-phosphonic acid (AMPA) occurred in 2007 only. The same sampling methods were used for glyphosate as for the other pesticides but a separate air sampling unit was used because the glyphosate compounds required a different analytical method than the multi-residue one used for the other pesticides and degradation products.

Most rain samples were composites of several precipitation events that occurred during the 7 day sampling periods. If more than 1 L of rainwater was collected during a sampling period, the total volume was recorded and a 1 L subsample was taken for analysis. The composite rain samples were collected in refrigerated glass carboys. To ensure reasonable analytical results, at least 75 mL of rainfall were required as a minimum sample size. This volume or greater was collected for all samples during both studies. The total amount of rainfall during the sample collection period was recorded by a rain gauge at each site, and the amount of rain from each event ranged from a trace amount to several centimeters (Figure 2). The weekly-integrated rainfall samples were collected over the same total period as the air samples. The mass of pesticide deposited by rain for the week (a weekly flux) was calculated by multiplying the concentration ( $\mu\text{g}/\text{m}^3$ ) by the rainfall amount (m) resulting in a mass per meter squared value. Weekly fluxes were summed to yield a cumulative, seasonal deposition flux.

The analytical methods used in 1995 and 2007 were very similar overall, but there were a few differences, primarily in the number of target analytes and in the applied reporting levels (RLs, see below). In 1995, the methods determined 47 analytes (Supplemental Data, Table S1) including four degradation products. The 2007 methods determined 76 analytes including 21 degradation products (Supplemental Data, Table S1). Thirty-four of the 1995 analytes were analyzed by the 2007 method, but of the 3 1995 degradation products, only CIAT was analyzed in 2007.

The air samples were analyzed using research methods [29, 31] adapted from the multi-residue pesticides in water method [32, 33] applied to the rain samples. Air sample components were individually extracted in a Soxhlet apparatus, and resultant extracts were passed through Florisil (1995) or C18 (2007) solid-phase extraction (SPE) columns. Rain samples were passed through C18 SPE columns and the pesticides were eluted using isopropanol (1995) or ethyl acetate (2007).

Air and rain sample extracts were analyzed by gas chromatography/mass spectrometry (GC/MS) using selected-ion monitoring with overall identical operational parameters between years. For 1995, applied rain RLs were set equivalent to the method detection limits (MDLs) estimated using the USEPA MDL procedure for the applied water method [32]. For 2007, rain RLs were set to twice the long-term MDLs, a value typically greater than the USEPA MDL [34]. Indeed, applied RLs in 2007 were at least twice those for 1995 (Supplemental Data, Table S1), and likely better reflect true RLs for the 1995 rain samples. Air method RLs were estimated based on

an 850 m<sup>3</sup> average air volume and using twice the corresponding water methods' MDLs [34], or from instrumental detection levels with consideration of method performance. Air and rain values below the MDLs and RLs were reported as estimated concentrations as described in Childress [34] and were used in data evaluations. Glyphosate and AMPA in air and rain samples were derivatized and analyzed by online SPE coupled to liquid chromatography/mass spectrometry [27].

Total (gas + particle) air concentrations are discussed except where noted. Although separate gas- and particle-phase air concentrations were determined in both years, an in-depth analysis of the observed gas-particle partitioning is beyond the scope of this present study. The mean concentrations and percent detection for each phase are presented in Supplemental Data, Tables S2 and S3. Mean air and rain values are of detections only, whereas the median values include the non-detections (ND).

Mississippi state-wide pesticide use for 1995 and 2007 was estimated using proprietary and US Department of Agriculture-National Agriculture Statistics Service (USDA-NASS) data (<http://www.pestmanagement.info/nass/>) using a method developed by Thelin and Stone [35]. "Local" pesticide use within a 20 km radius around each sampling site also was estimated from the state-wide use data, application rates, and mapped agricultural land within the 20 km radius area, in a manner similar to Nakagaki and Wolock [36]. The "local-use" estimates were examined to see if they would provide a better correlation than the state-wide estimates between use and the observed occurrences, concentrations, and deposition fluxes of the detected pesticides.

## **RESULTS**

### *Rainfall patterns in 1995 and 2007*

Figure 2 shows the weekly rainfall amounts and cumulative deposition from April through mid-September for both 1995 and 2007. Approximately 20% more rain fell in 1995 than in 2007. In 1995, the frequency and amounts of rainfall were both greater in the early months (April and May) of the growing season when preplant/pre-emergence herbicides were applied. In 2007, most of the rainfall events occurred in June after most of the herbicide had been applied. Accordingly, detection frequencies, concentrations, and deposition amounts for



most herbicides might be expected to be greater in 1995 than in 2007.

Sampling rainfall does not offer the continuous data record of atmospheric pesticide burden that air sampling does, but it does provide a direct measure of the atmospheric occurrence of a pesticide and the amount redeposited back to the earth's surface, including to unintended areas. Sampling rainfall also provides insight into the contribution of wet deposition to pesticide loads observed in runoff and surface waters.

#### *Summary of 1995 observations*

Thirty-two compounds were detected in the air or the rain in 1995 (Supplemental Data, Table S2). Twenty of these were common to both matrices (Figure 3A). Six were only detected in rain (alachlor, prometon, simazine, linuron, terbacil, and parathion) and 5 others were only detected in air (4,4'-DDE,  $\alpha$ -HCH,  $\gamma$ -HCH (lindane), metribuzin, and butylate). Eleven compounds were detected in 50% or more of all the rain or air samples collected in 1995 (Supplemental Data, Table S2). Seven of these (atrazine, metolachlor, molinate, methyl parathion, propanil, pendimethalin, and trifluralin) were detected in both air and rain at  $\geq 50\%$  frequency. The atrazine (and propazine) degradation product 2-chloro-4-isopropylamino-6-amino-*s*-triazine (CIAT; often called deethylatrazine) was detected in 71% of the air samples, but only in 38% of the rain samples. 2-Chloro-6-ethylamino-4-amino-*s*-triazine (CEAT), a degradation product of atrazine, cyanazine, simazine, and terbuthylazine (often called deisopropylatrazine) was detected in 38% of air samples, but was not analyzed in rain samples because of poor SPE collection efficiency.

DDE was detected in 100% of the gas-phase and 52% of the particle-phase air samples, but not in any rain sample. The mean DDE concentration ( $0.62 \text{ ng/m}^3$ ) was approximately 7 times greater than in ambient air samples from Louisiana and Texas, but less than samples taken 40 cm above the soil from those states and Alabama [37]. The occurrence of DDE was indicative of volatilization losses from past use of DDT. Nineteen of the 28 detected parent compounds were reportedly used in Mississippi and 4 others that were not detected had reported use. Nine pesticides had no reported use but were detected in either the air, rain, or both (Supplemental Data, Table S2).

#### *Summary of 2007 observations*

Thirty-seven compounds were detected in the air or rain samples in 2007 (Supplemental Data, Table S3). As was the case in 1995, 20 of these were present in both air and rain (Figure 3B). Four compounds were only detected in rain (the oxygen analog [OA] of methyl parathion, acetochlor, dichlorvos, and prometryn), and 13

compounds were only detected in air. Fourteen compounds were detected in 50% or more of the samples but only 5 of these (atrazine, glyphosate, AMPA, metolachlor/S-metolachlor, and propanil) were detected in 50% or more of the rain samples (Supplemental Data, Table S3). Nineteen of the 27 detected parent pesticides had reported state-wide use and 4 others that were analyzed for had reported use but were not detected. Seven had no reported use, but were detected in the air, rain, or both (Supplemental Data, Table S3). Cyanazine, which was detected in 29% of air and 56% of rain samples in 1995, was not detected in any 2007 air or rain samples; a clear reflection of its use cancellation in 1999. Molinate was also detected in both air and rain in 2007 even though its voluntary phase out began 4 years earlier. These detections were possibly the result of unreported use of stored molinate reserves.

Six compounds (CEAT, DDE,  $\alpha$ -HCH,  $\gamma$ -HCH, linuron, and parathion) were not analyzed for in 2007 because of programmatic changes to the GC/MS target analyte list. A variety of compounds (10 insecticides, 4 herbicides, and 5 fungicides) were added to the 2007 analyses including the herbicide glyphosate and one of its major degradation products AMPA, the insecticides fipronil and dicotophos, and the fungicide isomers *cis*- and *trans*-propiconazole. Glyphosate was the predominant new herbicide detected in both air (86%) and rain (77%); *cis*- and *trans*-propiconazole were detected between 30% and 50% in both air and rain. Use of many of these compounds either began or increased dramatically between collection years primarily because of the increased planting of GM crops or in response to the detection of soybean rust that began in 2005.

## **DISCUSSION**

### *Seasonal patterns*

Pesticide compounds were detected in both air and rain beginning in April 1995 and 2007 coinciding with the planting season. The sampling in 2007, however, started approximately two weeks later than in 1995 for both matrices.

*Herbicides:* Herbicides in air and rain were detected primarily in May of both years at nearly equivalent concentrations. Concentrations for many herbicides peaked around mid-May and then declined. By July concentrations were generally very low or not detectable. Figures 4A and B show the weekly concentrations in air for 4 of the most frequently detected herbicides (atrazine, metolachlor, pendimethalin, and propanil) in relation to

the sum of the other herbicides detected in 1995 and 2007, respectively. Figure 4B includes glyphosate that was not analyzed in 1995. These 4 (or 5) herbicides were detected in greater than 50% of the air samples.

In 1995, the "Other Herbicides" category dominated the concentrations for most air samples with three herbicides contributing to the weekly totals most frequently and with the highest concentrations. Trifluralin was detected in every sample with a median and mean concentration of detections of 0.81 ng/m<sup>3</sup> and 1.69 ng/m<sup>3</sup>, respectively. Molinate (62%, 0.13 and 1.00 ng/m<sup>3</sup>) and thiobencarb (38%, ND, 5.91 ng/m<sup>3</sup>), both rice herbicides, accounted for much of the rest.

In 2007, the "Other Herbicides" category made only minimal contributions to the total weekly observed air concentrations (Figure 4B). Trifluralin was again detected in nearly every sample (95%), but its median and mean concentrations were approximately 20 times less than in 1995 (Supplemental Data, Tables S2 and S3), similar to the nearly 20 fold decrease in reported use within 20 km of the sampling site. No molinate use was reported in 2007, yet it was detected in 24% of the air samples with a mean concentration only slightly less than in 1995. Conversely, thiobencarb use within 20 km of the sampling site was approximately 4 times higher than in 1995, yet the detection frequency (14%) and mean concentration (0.26 ng/m<sup>3</sup>) were both much lower than in 2007. That molinate concentrations in 2007 were not substantially different than in 1995 was likely due to the 2007 site being near more rice fields than the 1995 site.

Pendimethalin and propanil were detected in nearly every air sample from April through June in both 1995 and 2007 and were usually present at concentrations greater than other compounds detected during the same period. Both herbicides are used on rice, but pendimethalin also is used on field corn, potatoes, cotton, soybeans, tobacco, peanuts and sunflowers. For pendimethalin, changes in detection frequencies (62% to 57%) and mean concentrations (2.00 to 2.58 ng/m<sup>3</sup>) between 1995 and 2007 were not substantial, even though 2007 estimates of state-wide and local use were 27 and 8 times lower, respectively, than in 1995 (Supplemental Data, Tables S2 and S3). In contrast, the propanil detection frequency increased from 57% to 81% and the mean concentration decreased from 1.36 to 0.97 ng/m<sup>3</sup>. While the reported state-wide use decreased by two-thirds, the local use decreased only 12% (Supplemental Data, Tables S2 and S3). The increased propanil detection frequency is most likely due to the 2007 site being nearer to more rice fields than the 1995 site.

Metolachlor detections decreased from 81% in 1995 to 67% in 2007, but the mean concentration increased

more than 3 times from 0.20 to 0.74 ng/m<sup>3</sup> even though estimated state-wide and local use decreased by approximately 30% and more than 75%, respectively. In 1995, the metolachlor formulation contained racemic metolachlor (50% *S*- and *R*-metolachlor enantiomers). Beginning in 1998, use of a new formulation, called *S*-metolachlor, that contained 88% of the more herbicidally-active *S*-enantiomer and 12% *R*-enantiomer began displacing use of racemic metolachlor with the goal of reducing overall metolachlor application rates by 35% ([http://www.thecre.com/quality/20020305\\_SyngentaLet2CRE\\_reCedar.html](http://www.thecre.com/quality/20020305_SyngentaLet2CRE_reCedar.html)). In 2007, applications of racemic and *S*-metolachlor formulations occurred (2007 use and detections shown as “Metol+*S*-Metolachlor” in Supplemental Data, Table S3). Preferential use of *S*-metolachlor in 2007 might account for a portion of the reduced use of Metol+*S*-Metolachlor compared to 1995. There are no substantial differences in the physical/chemical properties (same vapor pressures, water solubilities, soil partition coefficients ([http://www.epa.gov/opp00001/chem\\_search/cleared\\_reviews/csr\\_PC-108801\\_28-Feb-02\\_234.pdf](http://www.epa.gov/opp00001/chem_search/cleared_reviews/csr_PC-108801_28-Feb-02_234.pdf)) between the *R*- and *S*-enantiomers to explain the 67% detection frequency relative to reduced use in 2007. Instead, closer proximity of the 2007 sampling site to application sites might explain both the detection frequency and the higher mean air concentration relative to 1995.

Atrazine use increased substantially state-wide from 1995 to 2007 due to over 3-fold increase in corn production and glyphosate use also increased due to the introduction of GM crops. While local glyphosate use increased proportionally to state-wide use in 2007, local atrazine use decreased by more than 36%. This discrepancy might be due to the manner by which local use was estimated – as a percent of state-wide use and agricultural acreage within the 20 km radius of the sampling site. It also may be due to the adoption of best management practices that included lower applications rates, the use of different formulations, or a change in weed control protocols for corn with the addition of glyphosate. The change in sampling location also may have contributed to the discrepancy because the distribution of crops receiving atrazine applications may have changed. The 2007 atrazine weekly air concentration pattern was markedly different from 1995 (Supplemental Data, Figures S1A and B). The first maximum 2007 concentration (1.24 ng/m<sup>3</sup>) occurred one week before the higher 1995 maximum (2.80 ng/m<sup>3</sup>), but planting was also 3 weeks earlier in 2007. The 2007 weekly concentrations over time,

however, did not drop off after mid-May as they did in 1995. In late July, there was a 5 week period where the air concentrations were very similar in level and pattern to what was observed during the normal application period in May 1995, with a secondary, late-season maximum concentration of 2.36 ng/m<sup>3</sup>. There are no known reasons for atrazine applications this late in the growing season. It is possible that these late season detections resulted from the entrainment of atrazine bound soil or pulverized plant tissue particles into the lower atmosphere during the harvesting process [38]. Since nearly all of the measured concentration was associated with the gas phase (Supplemental Data, Table S3 and Figure S1B), a likely source would be the volatilization of atrazine residues on the disturbed soil and pulverized plant tissue during the harvesting process. This late-season atrazine air concentration pattern has not been previously reported.

The 2007 weekly air concentration pattern for glyphosate (Figure 4B) was similar to those of other commonly detected herbicides in both 1995 and 2007 in that the highest concentrations occurred in April and May. However, there were detectable concentrations of glyphosate over the entire growing season, which is consistent with how glyphosate is used on GM crops, including for post-emergent weed control throughout the growing season.

Nine herbicides were detected in 50% or more of the rain samples in 1995 and only 3 of these (atrazine, metolachlor, and propanil) were detected in 50% or more of the 2007 rain samples, along with glyphosate (Supplemental Data, Tables S2 and S3). Detection patterns in rain were similar to those seen in corresponding air samples during both years for the more frequently detected herbicides, with the highest concentrations and most frequent rain detections occurring in April and May (Supplemental Data, Figures S2A–C). Both years also had very similar concentration ranges in rain even though detection frequencies and state-wide and local use were generally lower in 2007.

Several compounds with no reported use in Mississippi were detected in air and/or rain in 1995 or 2007. One is the herbicide DCPA, which was detected in air (24% in 1995; 95% in 2007) and rain ( $\leq 15\%$  both years) at very low concentrations (Supplemental Data, Tables S2 and S3). The combination of low or no use, variable detection frequencies, and very low concentrations might be an indicator of regional and long-range transport of pesticides into the study area, as observed by others for DCPA [39, 40]. The presence of DCPA in most air samples in 2007, coupled with very low concentrations, might also be a result of non-agricultural uses or use on

low acreage specialty crops within the sampling area.

*Insecticides:* Insecticides generally are not applied prophylactically like the pre-emergence herbicides, but as needed to combat specific insect infestations. Insecticide application rates and the overall mass used during a growing season are often considerably less than for most herbicides [11] (Supplemental Data, Tables S2 and S3).

The greatest insecticide air concentrations (Figure 5) during both years occurred later in the growing season than herbicides (Figure 4). Figure 5A shows 1995 weekly air concentrations for the three most frequently detected insecticides, and the sum of the other insecticides that were only detected once or twice in air. Methyl parathion was the most frequently detected (71%) insecticide in 1995, and it had the highest air concentrations of any compound during both study years with a mean concentration of  $14.2 \text{ ng/m}^3$ . It was heavily used (160,000 kg), primarily on cotton [41] in Mississippi. By 2007, methyl parathion's reported state-wide use dropped nearly 20 fold to 7,900 kg and as a result its detection frequency decreased to 29%. The methyl parathion concentrations (Figure 5B) were also considerably lower except during the week of 4 September, which had the greatest measured concentration ( $25.4 \text{ ng/m}^3$ ) of any detected compound in 2007 and exceeded all but 2 of the 1995 air concentrations. Excluding this value, the 2007 mean concentration of detections is  $0.47 \text{ ng/m}^3$ . Most weekly methyl parathion air concentrations were similar to ambient concentrations for California sites, and the timing of peak concentrations compares well with that previously observed for studies in Stoneville, MS in the 1970s, although our maximum concentrations were 2 to 100 times less (data summarized in [42]). Parathion (ethyl) had the highest reported use of all of the targeted insecticides, but was not detected in air and only detected once in rain. This is consistent with a short atmospheric lifetime [43]. The parathion OA degradation product was not measured in these 2 studies.

Chlorpyrifos was the second most frequently detected (38%) insecticide in 1995 air with a mean concentration of  $1.38 \text{ ng/m}^3$ . Malathion was detected less frequently (29%) than chlorpyrifos, but its state-wide and local uses were greater (Supplemental Data, Table S2), as was its mean concentration ( $3.64 \text{ ng/m}^3$ ), with some of the weekly values being similar in magnitude to those of methyl parathion. In 2007, the reported state-wide uses for chlorpyrifos and malathion were considerably less than in 1995 with both having no estimated local use. In

fact, the reported state-wide chlorpyrifos use dropped from over 19,000 kg in 1995 to 2 kg in 2007, yet the detection frequency in air increased to 95%, and its mean air concentration was only slightly less than in 1995. The mean concentration from mid-June to mid-September was  $1.23 \text{ ng/m}^3$ , nearly equal to that observed during the same time period in 1995. This indicates that the 2007 chlorpyrifos atmospheric burden was likely from a combination of local and regional sources [44]. The apparent discrepancy between the reported 2007 chlorpyrifos use and detection frequency also could be due to the way that the use was estimated. Thelin and Stone [35] reports that both low and high state-wide use values were based on harvested crop acreage and were estimated when data were unavailable. The low and high use estimates generally agreed well with other published values for selected pesticides, crops, and years, but not for some compounds. Chlorpyrifos is one compound where the low estimate of 2 kg (the value reported by the USDA-NASS) was very different from the high estimate of  $>14,000 \text{ kg}$ . This illustrates the uncertainty associated with comparing detection frequencies and observed concentrations to reported use data.

In 2007, the “Other Insecticides” category was mainly comprised of 4 pyrethroid insecticides and dicotophos. Both *cis*- and *trans*-permethrin detections were nearly 60%, with mean concentrations of approximately  $0.05 \text{ ng/m}^3$  each. Permethrin, however, had essentially no reported state-wide or local use.  $\lambda$ -Cyhalothrin had the highest reported state-wide use of the pyrethroid insecticides (8,237 kg), and was detected in 43% of the samples with a mean concentration of only  $0.07 \text{ ng/m}^3$ . These low concentrations are consistent with how insecticides, in general, and pyrethroid insecticides specifically, are applied with rates on the order of grams per hectare instead of kilograms per hectare used with some herbicides. They are also reflective of the low volatility of pyrethroid insecticides. Dicotophos and acephate (not measured) were the two primary insecticides used on cotton in 2007 that displaced methyl parathion use (<http://quickstats.nass.usda.gov/results/D33D31E3-1111-31A9-ADA2-5C075AC0C56E>; [45]). The detection frequency of dicotophos in air (24%) was similar to malathion, but its mean concentration was approximately one-third less ( $0.23 \text{ ng/m}^3$ ), even though its state-wide and local uses were at least 4 times greater than malathion.

The insecticide fipronil was not detected in air, but 2 of its degradation products, fipronil sulfide (62%) and desulfinyl fipronil (71%), were frequently detected at low levels ( $< 0.012 \text{ ng/m}^3$ , Supplemental Data, Table S3). Neither fipronil nor any targeted degradation products were detected in rain.

Concentrations in rain generally paralleled their counterparts in air for the most frequently detected insecticides (Supplemental Data, Figures S3A–C). Methyl parathion was detected in 69% of the rain samples (comparable to air). It was detected almost continuously from mid-May through July, 1995. It had a high mean concentration (3.04  $\mu\text{g/L}$ ) due to 2 very high sample concentrations (Supplemental Data, Figure S3A). The highest concentration in rain (22.9  $\mu\text{g/L}$ ;  $\sim 7$  cm of rainfall), which occurred in late June, did not correspond with the highest air concentrations, which occurred in August. However, the lack of rain over nearly the entire month of August (Figure 2) may have contributed to the high air concentration. Washout and rainout are very efficient mechanisms for removing airborne pesticide residues [14] and the greater the time between rainfall events, the greater the potential for more pesticides to become airborne through application drift, volatilization, or sorbed to soil particles entrained into the lower atmosphere by wind. In 2007, methyl parathion was only detected twice in rain in late August and early September at concentrations similar to those in 1995. All the other insecticides were detected in less than 40% of the 2007 rain samples. Malathion (Supplemental Data, Figure S3B) also was detected in only 2 samples in 2007 with nearly the same mean concentration as methyl parathion (0.71 and 0.75  $\mu\text{g/L}$ , respectively). This is consistent with the decrease in reported use. Although the rain detection frequencies of chlorpyrifos (38%) were the same in both years, its 2007 mean concentration (0.005  $\mu\text{g/L}$ ) was approximately 3 times less than in 1995 (0.017  $\mu\text{g/L}$ ). The "Other Insecticides" – the four pyrethroid insecticides and dicotophos – were not detected in any rain sample.

#### *Seasonal deposition by rain*

Detection frequencies and concentrations indicate how often a pesticide occurred in rain and what the potential exposure might be to non-target organisms, but the significance of these occurrences and concentrations also depends on the actual mass deposited by rainfall. The total seasonal rain deposition or flux ( $\mu\text{g/m}^2$ ) for each pesticide and degradation product is given in Supplemental Data, Tables S2 and S3.

The total pesticide flux in 1995 (1,981  $\mu\text{g/m}^2$ ) was 4.5 times greater than in 2007 (437  $\mu\text{g/m}^2$ ). Pesticide deposition was dominated by methyl parathion which comprised 88% of the total pesticide flux (Figure 6). Herbicides contributed only 11% (210  $\mu\text{g/m}^2$ ) of the total pesticide flux in 1995. Nearly all ( $>98\%$ ) the methyl



parathion deposition came from only two rain events and may have been a result of drift from nearby applications. Calculating an “adjusted” flux that does not include these two outlying values may provide a more realistic assessment of ambient conditions and better show the deposition contributions of the other pesticides.

Under this adjustment, the total insecticide deposition decreases to  $64 \mu\text{g}/\text{m}^2$ , 23% of the total deposition. The adjusted methyl parathion contribution ( $30 \mu\text{g}/\text{m}^2$ ) still contributes nearly one-half of the total insecticide deposition. The herbicide contribution to the adjusted total deposition increases to 76% with propanil and atrazine contributing over one-half (51%) of the total herbicide flux. Pendimethalin, metolachlor, and molinate together account for 33% of the total adjusted herbicide flux, with 11 others contributing the remaining 16%. Carbofuran, which was detected in only one-quarter of the rain samples, accounted for 25% of the adjusted insecticide flux, while chlorpyrifos and malathion accounted for 10%. Although detected only twice, azinphos methyl accounted for 6.5% of the adjusted total insecticide flux; nearly the same as more commonly detected malathion. The remaining 4 detected insecticides accounted for 5.7% of the adjusted total insecticide flux.

In 2007, herbicides contributed  $235 \mu\text{g}/\text{m}^2$  (54%) to the total pesticide deposition flux, a slight flux increase from 1995. Glyphosate accounted for 55% of the total herbicide flux, which was not surprising considering that more than 2 million kilograms were applied state-wide in 2007; 3 times that of atrazine and approximately one-third of the total reported applications for the other measured herbicides. The same 5 herbicides (propanil, atrazine, metolachlor, pendimethalin, and molinate) that contributed most to the overall herbicide flux in 1995 also contributed the most in 2007. Glyphosate was the exception because it was not analyzed for in 1995. It was likely present in some of the 1995 air and rain samples, but probably at lower concentrations than seen in 2007. Based on the 1995 reported use of glyphosate (147,000 kg state-wide and 4,751 kg within 20 km), and the ratio of the 2007 use to flux amount, glyphosate could have added an estimated 3% ( $\sim 7 \mu\text{g}/\text{m}^2$ ) to the 1995 total herbicide flux.

The introduction of glyphosate-resistant (GR) crops resulted in several significant changes in agriculture as well as in herbicide use. In some respects it has simplified weed management with glyphosate applications on GR soybean almost completely replacing the use of other herbicides [25]. In addition, because glyphosate has minimal, if any residual herbicide activity, some growers delay applications until several weeks after planting instead of the usual preplant or preemergence weed control [25] with the goal of reducing herbicide application costs and eliminating/reducing multiple applications. With the reliance on glyphosate for weed control in GR

crops, some weeds have evolved resistance to glyphosate [24, 46, 47] and there is now a need to diversify and rotate applications using herbicides with different modes of action [48].

In 2007, the atrazine flux increased by 36% relative to 1995, consistent with its state-wide increased use (although the estimated local use of atrazine decreased from 1995 to 2007). The propanil flux was approximately one-quarter of the 1995 value, also consistent with its decreased state-wide use. Eleven other herbicides were detected in 2007, but their total flux was less than one-half of that in 1995, with metolachlor, molinate, and pendimethalin contributing 95% of the “Other Herbicides” flux.

Insecticides contributed 33% ( $146 \mu\text{g}/\text{m}^2$ ) to the 2007 total pesticide flux. Methyl parathion (47%) and malathion (51%) dominated the insecticide portion, but all of their contributions (and almost one-half of the chlorpyrifos flux) came from just 2 large rainfall events in late August and early September that constituted nearly 25% of the total April-September rainfall (Supplemental Data, Figures S3A–C). Chlorpyrifos, which was detected predominantly in the gas-phase in nearly every 2007 air sample, was only detected in 5 rain samples and contributed less than 1% to the total insecticide flux. The low occurrence of chlorpyrifos in rain is not unexpected because its Henry’s law constant is approximately 50 and 500 times greater than methyl parathion and malathion, respectively [14]. Other insecticides like carbaryl, dichlorvos, and dicrotophos contributed only 2% to the 2007 total insecticide flux. Fungicides were only analyzed in 2007 and accounted for less than 2% of the total seasonal pesticide flux, all of it from *cis*- and *trans*-propiconazole.

Three degradation products (CIAT, DDE, and  $\alpha$ -HCH) were analyzed in 1995 but only CIAT was detected in rain, and it accounted for <1% of the total pesticide flux. The most commonly detected degradation products in 2007 were AMPA, CIAT, 3,4-dichloroaniline (a propanil degradation product), the OAs of chlorpyrifos, malathion, and methyl parathion, and two degradation products of fipronil – fipronil sulfide and desulfinyl fipronil. AMPA was the most frequently detected degradation product in both air (86%) and rain (77%) in 2007 and had the highest flux relative to the parent pesticide (AMPA flux was ~20% of the glyphosate flux). Both malathion OA and methyl parathion OA were detected in less than 25% of the rain samples, with fluxes approximately 10% of the parent compounds (Supplemental Data, Table S3). CIAT was detected in 46% of the rain samples and its seasonal

flux was approximately 13% of its presumed parent, atrazine. Although propanil was detected in more than 50% of the rain samples, its degradation product, 3,4-dichloroaniline, was detected in only 2 samples and contributed approximately 0.5% to the total seasonal flux.

*Reasons for observed differences between years*

The 1995 sampling site was somewhat removed from active fields, so the concentrations in air and rain would have reflected a more well-mixed atmosphere and been more indicative of regional concentrations. In spite of this, the methyl parathion concentrations in both rain and air were the highest measured for any compound during both years. These very high values indicate that use in the surrounding area was high, that off site drift was excessive, and that it rained shortly after nearby applications. The concentrations of the other detected insecticides and herbicides in 1995 were variable, but none were present in the very high concentrations measured for methyl parathion.

The 2007 sampling site location, adjacent to an active soybean field, may explain why the metolachlor detection frequencies and concentrations in both air and rain were similar to those in 1995 even though the reported use decreased by nearly one-half. The atrazine concentrations were consistent and comparable to those in 1995 even though the use increased nearly 3 fold. These results are indicative of a regional source. The 2007 methyl parathion concentrations were generally much lower than in 1995, but there was one relatively high air and rain concentration in early September that may have been from applications near the sampling site. It also may have been from regional transport as methyl parathion is extensively used on cotton, but there was no cotton grown in the immediate area of the sampling site. The extremely elevated concentrations and flux of methyl parathion in 1995 suggest a substantial local source.

The occurrence of pesticides in rain is dependent upon rainfall. The timing and amount of rain, relative to application, is an important determinant for what, how much, and when a pesticide is detected. Pesticide detection frequencies in rainfall are a function of the number of rainfall events that occur during the study period. The number of rainfall events can either increase or decrease the detection frequency. As an example, if only a few rainfall events occur during the sampling period, but all of them occur during the peak application time, then the detection frequencies could be very high. Conversely, if many rainfall events occur outside application periods, then the detection frequencies could be very low.

Low detection frequencies do not necessarily mean that the resulting concentrations or deposition amounts are insignificant. The last 2 rain events of the 2007 season occurred in August and September and accounted for nearly 25% of the total seasonal rainfall amount. These 2 rain events contributed nearly one-half of the seasonal deposition for chlorpyrifos and all of the deposition for malathion and methyl parathion (Supplemental Data, Figures S3A–C). These results show that 1 or 2 high concentrations in sizable rainfall events can add significantly to the seasonal total depositional loads.

A wide variety of pesticides and their degradation products were detected in both air and rain, and many of the same compounds were detected in both 1995 and 2007. Atrazine, metolachlor, and propanil were the most frequently detected pesticides in both air and rain during both years. The most noticeable change from 1995 to 2007 was the decrease in the annual amounts of pesticides used with the exception of glyphosate, atrazine, and a few others with minor use. The decrease in overall use generally resulted in decreased detection frequencies in air and rain, but those pesticides with the highest use generally were detected most frequently. The relationship between use and detection frequency was not a simple one, however, and the decrease in detections was generally not proportional to the decrease in use. In some cases, as with propanil and chlorpyrifos, detections actually increased with a supposed substantial decrease in use. In addition, some pesticides, like DCPA, were detected yet had no reported use while others had reported use but were not detected. Local use estimates helped explain some of these variations, but as with state-wide use, they did not consistently explain the variations.

The most compelling reasons why differences in pesticide occurrence and concentration in air and rain from 1995 to 2007 were seen are changes in use that include regulatory restrictions and replacement by other or newer compounds; changes in sampling site location with different cropping patterns surrounding the site; and variability in the timing, intensity, and amount of rainfall. It is more likely, however, that the observed differences were due to a combination of these factors. Although detection frequencies and the seasonal median concentrations were lower in 2007, it was difficult to ascertain whether pesticides came from local, regional, or long-range transport. To better understand the ecological and human health implications of the wide variety of pesticides detected in the air and rain, more detailed information on pesticide use is needed along with more

extensive air and rain sampling networks.

## **SUPPLEMENTAL DATA**

**Tables S1–S3.** (40 KB XLS)

**Figures S1–S3.** (490 KB PDF).

*Acknowledgment*—We thank M. Manning and ?? for sample collection assistance, F. Wiebe and R. ReVello for analysis of the 1995 and 2007 air samples, respectively, and additional NWQL staff that prepared and analyzed the rain samples from both studies. Financial support was provided by the USGS National Water Quality Assessment Program, the Toxic Substances Hydrology Program, and the National Water Quality Laboratory.

*Disclaimer*—Any use of trade, product, or firm names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

## REFERENCES

1. Glotfelty DE, Leech MM, Jersey J, Taylor AW. 1989. Volatilization and wind erosion of soil surface applied atrazine, simazine, alachlor, and toxaphene. *Journal of Agricultural and Food Chemistry* 37:546-551.
2. Majewski MS. 1999. Micrometeorological methods for measuring the post-application volatilization losses of pesticides. *Water, Air, and Soil Pollution* 115:83-113.
3. Bidleman TF, Leone A, Wong F, Vliet LV, Szeto S, Ripley BD. 2006. Emission of legacy chlorinated pesticides from agricultural and orchard soils in British Columbia, Canada. *Environmental Toxicology and Chemistry* 25:1448-1457.
4. Majewski MS. 1991. Sources, movement, and fate of airborne pesticides. In Frehse H, ed, *Pesticide chemistry; Advances in international research, development, and legislation, Proceedings of the seventh international congress of pesticide chemistry (IUPAC), Hamburg, 1990*. Verlagsgesellschaft mbH, Hamburg, Germany, pp 307-317.
5. Prueger JH, Gish TJ, McConnell LL, Mckee LG, Hatfield JL, Kustas WP. 2005. Solar radiation, relative humidity, and soil water effects on metolachlor volatilization. *Environmental Science and Technology* 39:5219-5226.
6. Ferrari F, Karpouzas DG, Trevisan M, Capri E. 2005. Measuring and predicting environmental concentrations of pesticides in air after application to paddy water systems. *Environmental Toxicology and Chemistry* 39:2968-2975.
7. Slinn WGN. 1977. Some approximations for the wet and dry removal of particles and gases from the atmosphere. *Water, Air, and Soil Pollution* 7:513-543.
8. Risebrough RW. 1990. Beyond long-range transport: A model of a global gas chromatographic system. In Kurtz DA, ed, *Long Range Transport of Pesticides*. Lewis Publishers, Inc., Chelsea, MI, pp 417-426.

9. Wania F, Mackay D. 1996. Tracking the distribution of persistent organic pollutants. *Environmental Science and Technology* 30:390A-396A.
10. Scheringer M, Wegmann F, Fenner K, Hungerbühler K. 2000. Investigation of the cold condensation of persistent organic pollutants with a global multimedia fate model. *Environmental Science and Technology* 34:1842-1850.
11. Gilliom RJ, Barbash JE, Crawford CG, Hamilton PA, Martin JD, Nakagaki N, Nowell LH, Scott JC, Stackelberg PE, Thelin GP, Wolock DM. 2006. Pesticides in the nation's streams and ground water, 1992-2001. Circular. US Geological Survey, Sacramento, CA.
12. Kuang ZH, McConnell LL, Torrents A, Meritt D, Tobash S. 2003. Atmospheric deposition of pesticides to an agricultural watershed of the Chesapeake Bay. *Journal of Environmental Quality* 32:1611-1622.
13. Vogel JR, Majewski MS, Capel PD. 2008. Pesticides in rain in four small agricultural watersheds. *Journal of Environmental Quality* 37:1101-1115.
14. Majewski MS, Capel PD. 1995. Pesticides in the Atmosphere: Distribution, trends, and governing factors, 1st ed. Ann Arbor Press, Inc., Chelsea, MI.
15. Waite DT, Bailey P, Sproule JF, Quiring DV, Chau DF, Bailey J, Cessna AJ. 2005. Atmospheric concentrations and dry and wet deposits of some herbicides currently used on the Canadian Prairies. *Chemosphere* 58:693-703.
16. Addison RF, Smith TG. 1974. Organochlorine residue levels in arctic ringed seals: Variation with age and sex. *Oikos* 25:335-337.
17. George JL, Frear EH. 1966. Pesticides in Antarctica. *Journal of Applied Ecology* 3:155-176.
18. Strachan WMJ, Huneault H. 1979. Polychlorinated biphenyls and organochlorine pesticides in Great Lakes precipitation. *J Great Lakes Res* 5:61-68.
19. Kutz FW, Yobs AR, Yang SC. 1976. National pesticide monitoring programs. In Lee Jr RE, ed, *Air pollution from pesticides and agricultural processes*. CRC Press, Cleveland, OH, pp 95-136.
20. Richards RP, Kramer JW, Baker DB, Krieger KA. 1987. Pesticides in rainwater in the northeastern United States. *Nature* 327:129-131.

21. Goolsby DA, Thurman EM, Pomes ML, Meyers MT, Battaglin WA. 1997. Herbicides and their metabolites in rainfall: Origin, transport, and deposition patterns across the midwestern and northeastern United States, 1990-1991. *Environmental Science and Technology* 31:1325-1333.
22. Hatfield J, Wesley C, Prueger J, Pfeiffer R. 1996. Herbicide and nitrate distribution in central Iowa rainfall. *Journal of Environmental Quality* 25:259-264.
23. Capel PD, Ma L, Wotzka PJ. 1998. Wet atmospheric deposition of pesticides in Minnesota, 1989-1994. 97-4026. Water-Resources Investigation Report. US Geological Survey, Mounds View, MN.
24. Benbrook CM. 2012. Impacts of genetically engineered crops on pesticide use in the U.S. – the first sixteen years. *Environmental Sciences Europe* 24:1-13.
25. Young BG. 2006. Changes in herbicide use patterns and production practices resulting from glyphosate-resistant crops. *Weed Technology* 20:301-307.
26. Coupe RH, Manning MA, Foreman WT, Goolsby DA, Majewski MS. 2000. Occurrence of pesticides in rain and air in urban and agricultural areas of Mississippi, April-September 1995. *Science of the Total Environment* 248:227-240.
27. Chang FC, Simcik MF, Capel PD. 2011. Occurrence and fate of the herbicide glyphosate and its degradate aminomethylphosphonic acid in the atmosphere. *Environmental Toxicology and Chemistry* 30:548–555.
28. McCarthy KA, Rose CE, Kalkhoff SJ. 2010. Environmental settings of the South Fork Iowa River Basin, Iowa, and the Bogue Phalia Basin, Mississippi, 2006–10. 2012-5021. Scientific Investigations Report. US Geological Survey, Reston, VA.
29. Foreman WT, Majewski MS, Goolsby DA, Wiebe FW, Coupe RH. 2000. Pesticides in the atmosphere of the Mississippi River Valley, Part II - Air. *Science of the Total Environment* 248:213-226.
30. Majewski MS, Foreman WT, Goolsby DA. 2000. Pesticides in the atmosphere of the Mississippi River Valley, Part I - Rain. *Science of the Total Environment* 248:201-212.



31. Mendez GO, Foreman WT, Morita A, Majewski MS. 2008. Water- and air-quality monitoring of Sweetwater Reservoir watershed, San Diego County, California - Phase one results, 2001-2003. 347. Data Series Report. US Geological Survey, Sacramento, CA.
32. Zaugg SD, Sandstrom MW, Smith SG, Fehlberg KM. 1995. Methods of analysis by the U.S. Geological Survey National Water Quality Laboratory-Determination of pesticides in water by C-18 solid-phase extraction and capillary-column gas chromatography/mass spectrometry with selected-ion monitoring. 95-181. Open-File Report. US Geological Survey, Denver, CO.
33. Sandstrom MW, Stoppel ME, Foreman WT, Schroeder MP. 2001. Methods of analysis by the U.S. Geological Survey National Water Quality Laboratory--Determination of moderate-use pesticides and selected degradates in water by C-18 solid-phase extraction and gas chromatography/mass spectrometry. 01-4098. Water-Resources Investigation Report. US Geological Survey, Denver, CO.
34. Childress CJO, Foreman WT, Connor BF, Maloney TJ. 1999. New reporting procedures based on long-term method detection levels and some considerations for interpretations of water-quality data provided by the US Geological Survey National Water Quality Laboratory. 99-193. Open-file report. US Geological Survey, Reston, VA.
35. Thelin GP, Stone WW, 2013. Estimation of annual agricultural pesticide use for counties of the conterminous United States, 1992–2009. US Geological Survey Scientific Investigations Report 2013-5009, 54 p.
36. Nakagaki N, Wolock DM. 2005. Estimation of agricultural pesticide use in drainage basins using land cover maps and county pesticide data. 2005-1188. Open-File Report. US Geological Survey, Sacramento, CA.
37. Bidleman TF, Leone A. 2004. Soil–air relationships for toxaphene in the southern United States. *Environmental Toxicology and Chemistry* 23:2337-2342.
38. Clymo, AS, Shin, JY, Holmén, BA. 2005. Herbicide sorption to fine particulate matter suspended downwind of agricultural operations: Field and laboratory investigations: *Environmental Science and Technology*, 39:421-430.

39. Landers DH, Simonich SM, Jaffe D, Geiser L, Campbell DH, Schwindt A, Schreck C, Kent M, Hafner W, Taylor HE, Hageman K, Usenko S, Ackerman L, Schrlau J, Rose N, Blett T, Erway MM. 2008. The fate, transport, and ecological impacts of airborne contaminants in western national parks (USA). EPA/600/R-07/138. U.S. Environmental Protection Agency, Office of Research and Development, NHEERL, Western Ecology Division, Corvallis, OR.
40. Mast MA, Foreman WT, Skaates SV. 2007. Current-use pesticides and organochlorine compounds in precipitation and lake sediment from two high-elevation national parks in the Western United States. *Archives of Environmental Contamination and Toxicology* 52:294-305.
41. U.S. Department of Agriculture NASS. 1996. Agricultural chemical usage, 1995. Field crops summary. Available from:  
[http://usda.mannlib.cornell.edu/reports/nassr/other/pcubb/agricultural\\_chemical\\_usage\\_field\\_crops\\_summary\\_03.25.96](http://usda.mannlib.cornell.edu/reports/nassr/other/pcubb/agricultural_chemical_usage_field_crops_summary_03.25.96).
42. California Environmental Protection Agency. 1999. Evaluation of methyl parathion as a toxic air contaminant. Part A, Environmental fate. Dept. of Pesticide Regulation, Environmental Monitoring and Pest Management Branch, Sacramento, CA. Available from:  
<http://cdm16254.contentdm.oclc.org/cdm/singleitem/collection/p178601ccp2/id/1657/rec/18>.
43. Winer AM, Atkinson R. 1990. Atmospheric reaction pathways and lifetimes for organophosphorus compounds. In Kurtz DA, ed, *Long Range Transport of Pesticides*. Lewis Publishing Co., Chelsea, MI, pp 115-126.
44. Muir DCG, Teixeira C, Wania F. 2004. Empirical and modeling evidence of regional atmospheric transport of current-use pesticides. *Environmental Toxicology and Chemistry* 23:2421-2432.
45. U.S. Environmental Protection Agency. 2002. Benefits assessment for methyl parathion use in cotton. Office of Prevention, Pesticides and Toxic Substances, Washington, DC, Document EPA-HQ-OPP-2003-0237-0008, p. 11, available from: <http://www.regulations.gov/#!documentDetail;D=EPA-HQ-OPP-2003-0237-0008>.

46. Johnson WG, Owen MDK, Kruger GR, Young BG, Shaw DR, Wilson RG, Wilcut JW, Jordan DL, Weller SC. 2009. US farmer awareness of glyphosate-resistant weeds and resistance management strategies. *Weed Technology* 23:308-312.
47. Preston C, Wakelin AM. 2008. Resistance to glyphosate from altered herbicide translocation patterns. *Pest Management Science* 64:372-376.
48. Prince JM, Shaw DR, Givens WA, Newman ME, Owen MDK, Weller SC, Young BG, Wilson RG, Jordan DL. 2012. Benchmark study: III. Survey on changing herbicide use patterns in glyphosate resistant cropping systems. *Weed Technology* 26:536-542.

### Figure legends

- Figure 1. Map showing the sampling locations in the Mississippi Delta agricultural region. The 1995 site was located near the city of Rolling Fork, MS and the 2007 site was located approximately 100 km north near the city of Pace, MS.
- Figure 2. Weekly and cumulative amount of rainfall from April through mid-September at the 1995 and 2007 sampling sites.
- Figure 3. Detection frequencies for those compounds detected in air or rain in 1995 (A) and 2007 (B).
- Figure 4. Weekly air concentrations for select herbicides in 1995 (A) and 2007 (B). Glyphosate determined in 2007 only.
- Figure 5. Weekly air concentrations for select insecticides in 1995 (A) and 2007 (B).
- Figure 6. Total seasonal deposition for the detected pesticides and degradation products in 1995 and 2007.

it




Figure 1

AC





Figure 2

A



B




Figure 3

Accepte





Figure 4



A



B




Figure 5




Figure 6