



This Issue

Views **25,620**

Citations **39**

Altmetric **956**

Comments **1**



PDF



FREE

Original Investigation

July 2016

Associations Between Cannabis Use and Physical Health Problems in Early Midlife

A Longitudinal Comparison of Persistent Cannabis vs Tobacco Users

Madeline H. Meier, PhD¹; Avshalom Caspi, PhD^{2,3,4}; Magdalena Cerdá, DrPH⁵; et al

[» Author Affiliations](#) | [Article Information](#)

JAMA Psychiatry. 2016;73(7):731-740. doi:10.1001/jamapsychiatry.2016.0637



Abstract

Importance After major policy changes in the United States, policymakers, health care professionals, and the general public seek information about whether recreational cannabis use is associated with physical health problems later in life.

Objective To test associations between cannabis use over 20 years and a variety of physical health indexes at early midlife.

Design, Setting, and Participants Participants belonged to a representative birth cohort of 1037 individuals born in Dunedin, New Zealand, in 1972 and 1973 and followed to age 38 years, with 95% retention (the Dunedin Multidisciplinary Health and Development Study). We tested whether cannabis use from ages 18 to 38 years was associated with physical health at age 38, even after controlling for tobacco use, childhood health, and childhood socioeconomic status. We also tested whether cannabis use from ages 26 to 38 years was associated with within-individual health decline using the same measures of health at both ages.

Exposures We assessed frequency of cannabis use and cannabis dependence at ages 18, 21, 26, 32, and 38 years.

Main Outcomes and Measures We obtained laboratory measures of physical health (periodontal health, lung function, systemic inflammation, and metabolic health), as well as self-reported physical health, at ages 26 and 38 years.

Results The 1037 study participants were 51.6% male (n = 535). Of these, 484 had ever used tobacco daily and 675 had ever used cannabis. Cannabis use was

associated with poorer periodontal health at age 38 years and within-individual decline in periodontal health from ages 26 to 38 years. For example, cannabis joint-years from ages 18 to 38 years was associated with poorer periodontal health at age 38 years, even after controlling for tobacco pack-years ($\beta = 0.12$; 95% CI, 0.05-0.18; $P < .001$). Additionally, cannabis joint-years from ages 26 to 38 years was associated with poorer periodontal health at age 38 years, even after accounting for periodontal health at age 26 years and tobacco pack-years ($\beta = 0.10$; 95% CI, 0.05-0.16; $P < .001$) However, cannabis use was unrelated to other physical health problems. Unlike cannabis use, tobacco use was associated with worse lung function, systemic inflammation, and metabolic health at age 38 years, as well as within-individual decline in health from ages 26 to 38 years.

Conclusions and Relevance Cannabis use for up to 20 years is associated with periodontal disease but is not associated with other physical health problems in early midlife.

Advertisement

Introduction

After major policy changes in the United States, policymakers, health care professionals, and the general public seek information about whether recreational cannabis use is associated with physical health problems later in life. Two recent reviews found that persistent cannabis use is associated with few physical health problems, with the possible exceptions being cardiovascular risks and bronchitis.^{1,2} However, firm conclusions cannot be drawn due to methodological shortcomings.³ Most studies⁴⁻¹¹ are cross-sectional or rely on self-reported health. These designs cannot resolve the temporal association between cannabis use and health, nor can they address the possibility that cannabis users may have biased perceptions of their health. Longitudinal studies with laboratory-based measures and physical examinations are needed.

Few longitudinal studies have characterized cannabis users' long-term health using objective, laboratory-based indexes and examinations (eTable 1 in the **Supplement**). Each study in eTable 1 in the **Supplement** focused on a single domain of physical health, providing an important but incomplete picture. In the present population-representative study of individuals followed up from birth to age 38 years, we tested associations between cannabis use over 20 years and multiple domains of physical health in early midlife. We selected the following health domains based on prior research,¹⁻³ demonstrated capacity to predict disease morbidity and mortality,¹²⁻¹⁴ and biological plausibility of an effect of cannabis by early midlife: periodontal health, lung function, systemic inflammation, and metabolic risk. First, we tested whether cannabis use from ages 18 to 38 years was associated with health at age 38 years. Second, we tested whether cannabis use from ages 26 to 38 years was associated with within-individual health decline using the same measures of health at both ages. To provide a benchmark for comparison, we also tested associations between tobacco use and physical health.

Key Points

Question Is persistent cannabis use for up to 20 years associated with physical health problems (periodontal health, lung function, systemic inflammation, and metabolic health) in early midlife?

Findings In this prospective, longitudinal study of a representative birth cohort of 1037 individuals, persistent cannabis use from ages 18 to 38 years was not associated with physical health problems at age 38 years with one exception: persistent cannabis use was statistically significantly associated with poor periodontal health.

Meaning Persistent cannabis use for up to 20 years is, for the most part, not associated with physical health problems in early midlife.

Methods

Participants

Participants were members of the Dunedin Multidisciplinary Health and Development Study of New Zealand, a longitudinal investigation of health and behavior in a representative birth cohort.¹⁵ Study members ($n=1037$), representing 91.0% of eligible births and 51.6% male, were all born in 1972-1973 in Dunedin, New Zealand. Of the 1037 study participants 51.6% were male ($n = 535$). They were eligible for the longitudinal study based on residence in the province at age 3 years and participation in the first follow-up at age 3 years. The cohort represents the full range of socioeconomic status (SES) in the general population of New Zealand's South Island and is primarily white.¹⁵ On adult health, the cohort matches the New Zealand National Health and Nutrition Survey (eg, body mass index, smoking status, and general practitioner visits).¹⁵ Assessments occurred at birth and ages 3, 5, 7, 9, 11, 13, 15, 18, 21, 26, 32, and, most recently, 38 years, when 95.4% ($n=961$) of the 1007 living study members took part. At each assessment phase, study members are brought to the Dunedin Multidisciplinary Health and Development Research Unit (Dunedin, New Zealand) for interviews and examinations. The Otago Ethics Committee (University of Otago, Dunedin, New Zealand) approved each phase of the study, and the study protocol was approved by the Otago Ethics Committee and the Duke University Institutional Review Board. Written informed consent was obtained from all participants.

Analyses were limited to 947 study members with laboratory health data at age 38 years because 46 study members were not seen at age 38 years, 30 were deceased, and 14 had field interviews that did not include laboratory measurements or examinations. There were no differences between those with and without health data at age 38 years on childhood health ($F=1.42$, $P=.23$), cigarettes smoked per day at age 18 years ($F=1.28$, $P=.26$), or frequency of cannabis use at

smoked per day at age 18 years ($F = 1.28, P = .26$), or frequency of cannabis use at age 18 years ($F = 2.85, P = .09$). **Table 1** lists characteristics of participants according to tobacco and cannabis exposure, including sex, childhood health,¹⁶ and childhood SES,¹⁷ which were available as covariates.

Tobacco Pack-Years

Cumulative tobacco exposure was calculated from the reported number of cigarettes smoked per day at each assessment divided by 20 and multiplied by the number of years smoked at that rate through age 38 years. One pack-year reflects the equivalent of 20 cigarettes a day for 1 year. Mean (SD) pack-years for those with age-38 health data was 6.17 (8.69). For analyses testing associations between pack-years from age 26 to 38 and change in physical health using the same measure of health at both ages, we estimated pack-years in the same way except that estimates represented cigarette use at ages 26 to 38 years (mean [SD], 3.30 [5.12] pack-years).

Cannabis Joint-Years

Pack-years, which combines information about smoking duration and intensity, is the most commonly used exposure in tobacco studies.¹⁸ We created a parallel variable that indexes cannabis smoking. Cumulative joint-years was estimated using self-reported frequency of cannabis use over the past year (0-365 days) at ages 18 to 38 years. One joint-year reflects the equivalent of daily cannabis use for 1 year. Mean (SD) joint-years between ages 18 to 38 years for those with age-38 health data was 1.99 (4.43). For analyses of health change from age 26 to 38 years, we estimated joint-years in the same way except that estimates represented cannabis use at ages 26 to 38 (mean [SD], 1.18 [3.00]).

Persistent Cannabis Dependence

Because prior reports from the Dunedin Study have characterized cannabis users in terms of persistent dependence over time,^{19,20} we also report this variable as our exposure. We assessed past-year dependence at ages 18 to 38 years with the

Diagnostic Interview Schedule^{21,22} according to *Diagnostic and Statistical Manual of Mental Disorders* criteria.^{23,24} Persistent dependence was defined as the number of study waves out of 5 at which a study member met criteria for dependence: never used cannabis at any study wave, used at least once between ages 18 to 38 but never diagnosed, diagnosed at 1 wave, diagnosed at 2 waves, and diagnosed at 3 or more waves. For analyses of health change from ages 26 to 38 years, we again defined persistent dependence as the number of study waves at which a study member met criteria for dependence but only used cannabis data from ages 26 to 38 years.

Physical Health at Age 38 Years

Physical examinations were conducted during the assessment day at age 38 years, with blood draws between 4:15 and 4:45 PM.²⁵ eTable 2 in the **Supplement** describes each health outcome: periodontal health, lung function, systemic inflammation, metabolic syndrome, waist circumference, high-density lipoprotein cholesterol level, triglyceride level, blood pressure (systolic and diastolic), glycated hemoglobin level, body mass index, and self-reported health. We report health outcomes scored as continuously distributed outcomes because continuous measures are more sensitive to variation than categorical measures. However, for clinical relevance, the eTables in the **Supplement** also list results for health outcomes scored as categorical clinical outcomes. Positively skewed continuous outcomes (combined attachment loss, inflammation, triglyceride level, and glycated hemoglobin level) were log-transformed before analysis.

Physical Health at Age 26 Years

The health measures at age 38 years were also administered at age 26 years using the same procedures, with 2 exceptions.²⁵ First, periodontal measurements were obtained using a half-mouth design.²⁶ Second, serum C-reactive protein level was assayed with a sensitivity level of 1 mg/L.²⁷ Due to this lower sensitivity, C-reactive protein levels in the top quintile of the distribution were designat-

ed as elevated.

Statistical Analysis

To test whether cannabis use was associated with poor health in early midlife, we tested the bivariate association between cannabis use from ages 18 to 38 years and health at age 38 years (**Table 2**, model 1) and subsequently added tobacco pack-years from ages 18 to 38 years as a covariate (**Table 2**, model 2). To test whether cannabis use from ages 26 to 38 years was associated with health decline using the same measure of health at both ages, we tested the bivariate association between cannabis use from ages 26 to 38 years and health at age 38 years (**Table 3**, model 1) and subsequently added health at age 26 years as a covariate (**Table 3**, model 2), followed by tobacco pack-years from ages 26 to 38 years as an additional covariate (**Table 3**, model 3). All analyses controlled for sex.

Statistical analyses tested associations of tobacco pack-years (a continuous variable), cannabis joint-years (a continuous variable), and cannabis dependence (a 5-level ordinal variable) with continuous and categorical health outcomes. We analyzed continuous outcomes using ordinary least squares regression to derive β coefficients and categorical outcomes using Poisson regression models to derive relative risks (RRs). We standardized continuous variables before conducting statistical tests. Therefore, β coefficients and RRs can be interpreted as the increase in risk of the outcome given a 1-SD increase in pack-years or joint-years. To aid interpretation of β coefficients and RRs associated with continuous pack-years and joint-years, we report unstandardized, sex-adjusted means for health outcomes as a function of tobacco and cannabis use, with study members grouped according to pack-years and joint-years in 5-year increments (**Table 2**).

Results

Tobacco Smoking and Health

Bivariate associations showed that tobacco pack-years was associated with worse health for 8 of the 12 health outcomes, namely, periodontal health, lung function, systemic inflammation, metabolic syndrome, high-density lipoprotein cholesterol level, triglyceride level, glycated hemoglobin level, and self-reported health (**Table 2**, model 1). Associations remained significant for all 8 of these health outcomes after controlling for cannabis joint-years (**Table 2**, model 2) and after additionally controlling for childhood health and SES (eTable 3, model 3 in the **Supplement**). Results were similar for clinically relevant, categorically scored health outcomes (eTable 3 in the **Supplement**). For example, 12.3% of individuals who never used tobacco daily had periodontal disease (≥ 1 site with attachment loss > 5 mm) compared with 52.9% of individuals with 15 or more pack-years (eTable 3 in the **Supplement**). Statistical tests showed that for every SD increase in pack-years (approximately 9 pack-years), the RR for periodontal disease increased by 1.63 ($P < .001$) (eTable 3 in the **Supplement**). Findings are consistent with prior research.^{26,28-36}

Cannabis Use and Health

Bivariate associations showed that cannabis joint-years was associated with worse health for 3 of 12 health outcomes, namely, periodontal health, lung function, and self-reported health (**Table 2**, model 1). Adverse associations remained significant for 2 outcomes (periodontal health and lung function) after controlling for tobacco pack-years (**Table 2**, model 2) and after additionally controlling for childhood health and SES (eTable 3, model 3 in the **Supplement**). However, poorer lung function (ratio of forced expiratory volume in the first second of expiration [FEV₁] to forced vital capacity [FVC]) (hereafter FEV₁/FVC) among cannabis users was probably not indicative of airway obstruction because joint-years (unlike tobacco pack-years) was unrelated to reduced FEV₁ (eTable 4 in the **Supplement**). Rather, reduced FEV₁/FVC among cannabis users was attributable to higher FVC values. It is unclear whether higher FVC values reflect better health.

Unlike tobacco, cannabis joint-years was associated with slightly smaller waist circumference and lower body mass index. Furthermore, after adjusting for tobacco pack-years (**Table 2**, model 2), associations emerged between joint-years and better high-density lipoprotein cholesterol, triglyceride, and glycated hemoglobin levels. However, joint-years was not associated with lower risk of metabolic syndrome. Results were similar for categorically scored health outcomes (eTable 3 in the **Supplement**).

Results for persistent cannabis dependence (as well as results for persistent regular cannabis use [eTable 5 in the **Supplement**]) were almost identical to those for joint-years. Bivariate associations showed that persistent dependence was associated with worse health for 3 of 12 outcomes, namely, periodontal health, lung function, and self-reported health. Associations remained significant for 1 of those 3 (periodontal health) after controlling for tobacco pack-years (**Table 2**, model 2) and after additionally controlling for childhood health and SES (eTable 3, model 3 in the **Supplement**). Results were similar for categorically scored health outcomes (eTable 3 in the **Supplement**). eTable 6 in the **Supplement** summarizes the aforementioned findings.

Periodontal health was the only aspect of health that showed a robust adverse association in analyses of both persistent dependence and joint-years. Post hoc analyses showed that cannabis users brushed and flossed less than others and were more likely to be alcohol dependent (eTable 7 in the **Supplement**). However, associations between cannabis use and poor periodontal health remained significant after controlling for tobacco pack-years, childhood health and SES, brushing and flossing, and alcohol dependence (eTable 8 in the **Supplement**).

The general lack of association between persistent cannabis use and poor physical health may be surprising. One explanation is that healthy youth select into cannabis use. Our test showed no correlation between cannabis use and childhood health (**Table 1**). Another explanation is that cannabis users may have

healthier adult lifestyles. Tests showed that cannabis use was not associated with more physical activity, with a diet of fruits and vegetables, or with less alcohol abuse (eTable 7 in the [Supplement](#)). The absence of associations between cannabis use and poor physical midlife health could not be attributed to better initial health, more physical activity, better diet, or less alcohol abuse.

Tobacco and Cannabis Use and Change in Health

Tobacco pack-years from ages 26 to 38 years was associated with worsening periodontal health, lung function, systemic inflammation, and metabolic health ([Table 3](#), model 2). For example, pack-years from ages 26 to 38 years was associated with increased risk of metabolic syndrome at age 38 years after accounting for metabolic syndrome at age 26 years (RR, 1.18; $P = .02$) ([Table 3](#), model 2). Tobacco users also self-reported worse health at age 38 years, and this association persisted after accounting for self-reported health at age 26 years ([Table 3](#), model 2).

Like tobacco use, cannabis use was associated with decline in periodontal health and lung function ([Table 3](#), model 2) even after accounting for tobacco pack-years from ages 26 to 38 years ([Table 3](#), model 3). Again, however, decline in the FEV₁/FVC was probably not attributable to airway obstruction because cannabis use was not robustly associated with decline in FEV₁ (eTable 9 in the [Supplement](#)). Cannabis use was not associated with deteriorating health in other domains. Results were similar for continuously scored and categorically scored health outcomes (eTable 10 in the [Supplement](#)).

Discussion

In general, our findings showed that cannabis use over 20 years was unrelated to health problems in early midlife. Across several domains of health (periodontal health, lung function, systemic inflammation, and metabolic health), clear evidence of an adverse association with cannabis use was apparent for only one

domain, namely, periodontal health. Cannabis use from ages 26 to 38 years was not associated with within-individual health decline during this 12-year period with the exception of periodontal health. By comparison, tobacco use was associated with worse periodontal health, lung function, systemic inflammation, high-density lipoprotein cholesterol levels, triglyceride levels, and glucose levels in early midlife, as well as health decline from ages 26 to 38 years.

Findings showed that cannabis use was associated with slightly better metabolic health (smaller waist circumference, lower body mass index, better lipid profiles, and improved glucose control). However, most of these associations emerged only after controlling for tobacco use. Effects were small but intriguing given similar reports from cross-sectional studies,^{4,8,9,37-39} as well as evidence that endocannabinoids appear to be involved in the regulation of metabolism.⁴⁰ Studies^{41,42} have shown that overweight patients who took rimonabant (a synthetic cannabinoid-1 receptor blocker) demonstrated reduced waist circumference and improved lipid profiles. However, it is unclear if and how recreational cannabis use (and plant-based cannabinoids) might affect metabolic health. Cannabinoid pharmacology is more complex than commonly believed,⁴³ and biological arguments can be made for cannabis-related worsening or improvement of metabolic health.^{9,44} The only other longitudinal study⁴⁵ to characterize cannabis users' metabolic health found no association, and our finding of a small association mainly emerged after controlling for tobacco use. Moreover, cannabis use was not associated with reduced risk of metabolic syndrome. Therefore, current evidence suggests that recreational cannabis use is unlikely to improve metabolic health in the general population.

In at least 2 instances, we found no association between cannabis use and poor health when we might have expected one. In the first instance, we found no association between cannabis use and reduced FEV₁ (eTable 4 in the **Supplement**), which is somewhat puzzling given that tobacco use is associated with reduced FEV₁.^{28,29} An association between cannabis use and reduced FEV₁ could emerge

with greater exposure to cannabis.²⁸ Nonetheless, given no evidence of reduced FEV₁ among cannabis users, our finding of lower FEV₁/FVC among cannabis users probably did not indicate airway obstruction. Rather, reduced FEV₁/FVC appeared to reflect cannabis users' slightly larger FVC. This association with larger FVC, also reported elsewhere,²⁸ is not understood. Overall, findings are consistent with a 2011 review concluding that there is little evidence that cannabis affects FEV₁ and airway obstruction.⁴⁶ In the second surprising instance, we found no association between cannabis use and cardiovascular risks (eg, high blood pressure and worse cholesterol levels), which may appear at odds with evidence that cannabis use increases risk for cardiovascular complications,⁴⁷⁻⁴⁹ even among young healthy individuals.⁵⁰ Our somewhat disparate findings are reconciled by evidence that cannabis-related cardiovascular complications are likely acute cannabis effects.^{38,45,47,49}

Although we found that cannabis users were generally no worse off than nonusers on almost all health indexes, they did have worse periodontal health. Cannabis use was associated with attachment loss, which can result in tooth loss.^{26,36} A similar association was observed for tobacco use, consistent with previous research.^{26,36,51} Tobacco's effect on periodontal disease is thought to be mediated through increased inflammation and vasoconstriction,⁵¹ which may or may not be the case for cannabis. Cannabis use was not associated with systemic inflammation here or elsewhere,^{30,39} but prior research has shown that cannabis use can induce vasoconstriction.^{52,53}

This study has limitations. First, cannabis joint-years was based on self-reports obtained at ages 18 to 38 years. Validation of cannabis use through laboratory measures could have helped detect cannabis users who denied use. However, underreporting due to reluctance to admit to illegal drug use is unlikely because study members, interviewed repeatedly over the course of their lives, have learned to trust our confidentiality guarantee. Second, disentangling cannabis use and tobacco use is challenging. In New Zealand, cannabis is not typically

mixed with tobacco,¹⁰ but most participants who used cannabis also smoked cigarettes. Although we controlled for tobacco use, imperfect control might bias results toward finding spurious associations between cannabis use and poor health. However, we note that all poor health outcomes, apart from periodontal disease, were unrelated to cannabis use. Third, our findings are based on a single New Zealand cohort who began using cannabis in the 1980s to 1990s. Although our findings are generally consistent with longitudinal studies^{2,54} of US samples (eTable 1 in the **Supplement**), tetrahydrocannabinol (the primary psychoactive ingredient in cannabis) content has increased since then.⁵⁴ If health associations are mediated by tetrahydrocannabinol, we may have underestimated the association between cannabis use and health. Fourth, our conclusions are limited to a specific set of health problems assessed in early midlife. Although this investigation is the most comprehensive study to date, cannabis use may be associated with health problems not studied here or with those that tend to emerge later in life, such as cancer. Fifth, we compared findings for cannabis against findings for tobacco. Our intent in doing so was to allay concerns that our study's methods might be unable to detect health problems. We acknowledge that participants acquired more tobacco pack-years than cannabis joint-years, with most cannabis users using daily for fewer than 5 years. Greater tobacco exposure may explain health decline associated with tobacco use but not cannabis use. If patterns of cannabis use shift and more users begin to use cannabis as they do tobacco (ie, multiple joints per day), cannabis-associated health problems might emerge. Sixth, our study cannot comment on the health effects of cannabis use in older adults or the safety of medical marijuana use in patients who are already unwell.

Conclusions

This study has a number of implications. First, cannabis use for up to 20 years is not associated with a specific set of physical health problems in early midlife. The sole exception is that cannabis use is associated with periodontal disease. Second, cannabis use for up to 20 years is not associated with net metabolic

Second, cannabis use for up to 20 years is not associated with net metabolic benefits (ie, lower rates of metabolic syndrome). Third, our results should be interpreted in the context of prior research showing that cannabis use is associated with accidents and injuries, bronchitis, acute cardiovascular events, and, possibly, infectious diseases and cancer, as well as poor psychosocial and mental health outcomes.^{1-3,19,20,46,55-58}

[Back to top](#)

Article Information

Submitted for Publication: November 25, 2015; final revision received January 26, 2016; accepted February 27, 2016.

Corresponding Author: Madeline H. Meier, PhD, Department of Psychology, Arizona State University, PO Box 871104, Tempe, AZ 85287-1104 (madeline.meier@asu.edu).

Published Online: June 1, 2016. doi:[10.1001/jamapsychiatry.2016.0637](https://doi.org/10.1001/jamapsychiatry.2016.0637).

Author Contributions: Dr Meier had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Meier, Caspi, Cerdá, Poulton, Thomson, Moffitt.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Meier, Caspi, Thomson.

Critical revision of the manuscript for important intellectual content: Meier, Cerdá, Hancox, Harrington, Houts, Poulton, Ramrakha, Thomson, Moffitt.

Statistical analysis: Meier, Caspi, Houts.

Obtained funding: Caspi, Poulton, Moffitt.

Administrative, technical, or material support: Harrington, Poulton, Ramrakha.

Study supervision: Cerdá, Poulton, Moffitt.

Conflict of Interest Disclosures: None reported.

Funding/Support: The Dunedin Multidisciplinary Health and Development Research Unit is supported by the New Zealand Health Research Council. This research was funded by grants R01AG032282 and R01AG048895 from the US National Institute on Aging, by grant MR/K00381X from the United Kingdom Medical Research Council, and by grant ES/M010309/1 from the Economic and Social Research Council. Additional support was provided by the Jacobs Foundation.

Role of the Funder/Sponsor: The funding agencies were not involved in the design or conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Previous Presentation: This study was presented at the 2016 Life History Research Society Meeting; May 25-28, 2016; Amsterdam, the Netherlands.

Additional Contributions: We thank study founder Phillip Silva, PhD, the Dunedin Multidisciplinary Health and Development Study of New Zealand members, and the Dunedin Multidisciplinary Health and Development Research Unit research staff (none received compensation).

References

1. Hall W. What has research over the past two decades revealed about the adverse health effects of recreational cannabis use? *Addiction*. 2015;110(1):19-35.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

2. Volkow ND, Baler RD, Compton WM, Weiss SRB. Adverse health effects of marijuana use. *N Engl J Med*. 2014;370(23):2219-2227.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

3. Gordon AJ, Conley JW, Gordon JM. Medical consequences of marijuana use: a review of current literature. *Curr Psychiatry Rep*. 2013;15(12):419.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

4. Le Strat Y, Le Foll B. Obesity and cannabis use: results from 2 representative national surveys. *Am J Epidemiol*. 2011;174(8):929-933.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

5. Baggio S, N'Goran AA, Deline S, et al. Patterns of cannabis use and prospective associations with health issues among young males. *Addiction*. 2014;109(6):937-945.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

6. Georgiades K, Boyle MH. Adolescent tobacco and cannabis use: young adult outcomes from the Ontario Child Health Study. *J Child Psychol Psychiatry*. 2007;48(7):724-731.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

7. Huang DY, Lanza HI, Anglin MD. Association between adolescent substance use and obesity in young adulthood: a group-based dual trajectory analysis. *Addict Behav*. 2013;38(11):2653-2660.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

8. Rajavashisth TB, Shaheen M, Norris KC, et al. Decreased prevalence of diabetes in marijuana users: cross-sectional data from the National Health and Nutrition Examination Survey (NHANES) III. *BMJ Open*. 2012;2(1):e000494. doi:10.1136/bmjopen-2011-000494.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

- 9.** Penner EA, Buettner H, Mittleman MA. The impact of marijuana use on glucose, insulin, and insulin resistance among US adults. *Am J Med.* 2013;126(7):583-589.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 10.** Aldington S, Williams M, Nowitz M, et al. Effects of cannabis on pulmonary structure, function and symptoms. *Thorax.* 2007;62(12):1058-1063.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 11.** Bechtold J, Simpson T, White HR, Pardini D. Chronic adolescent marijuana use as a risk factor for physical and mental health problems in young adult men. *Psychol Addict Behav.* 2015;29(3):552-563.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 12.** Danesh J, Whincup P, Walker M, et al. Low grade inflammation and coronary heart disease: prospective study and updated meta-analyses. *BMJ.* 2000;321(7255):199-204.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 13.** Eckel RH, Alberti KG, Grundy SM, Zimmet PZ. The metabolic syndrome. *Lancet.* 2010;375(9710):181-183.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 14.** Rasmussen F, Taylor DR, Flannery EM, et al. Risk factors for airway remodeling in asthma manifested by a low postbronchodilator FEV1/vital capacity ratio: a longitudinal population study from childhood to adulthood. *Am J Respir Crit Care Med.* 2002;165(11):1480-1488.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 15.** Poulton R, Moffitt TE, Silva PA. The Dunedin Multidisciplinary Health and Development Study: overview of the first 40 years, with an eye to the future. *Soc Psychiatry Psychiatr Epidemiol.* 2015;50(5):679-693.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)

- 16.** Belsky DW, Caspi A, Israel S, Blumenthal JA, Poulton R, Moffitt TE. Cardiorespiratory fitness and cognitive function in midlife: neuroprotection or neuroselection? *Ann Neurol.* 2015;77(4):607-617.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 17.** Elley WB, Irving JC. Revised socio-economic index for New Zealand. *N Z J Educ Stud.* 1976;11(1):25-36.
[Google Scholar](#)
- 18.** Thomas DC. Invited commentary: is it time to retire the "pack-years" variable? maybe not! *Am J Epidemiol.* 2014;179(3):299-302.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 19.** Meier MH, Caspi A, Ambler A, et al. Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proc Natl Acad Sci U S A.* 2012;109(40):E2657-E2664. doi:[10.1073/pnas.1206820109](https://doi.org/10.1073/pnas.1206820109).
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 20.** Cerdá M, Moffitt TE, Meier MH, et al. Persistent cannabis dependence and alcohol dependence represent comparable risks for midlife economic and social problems: a longitudinal cohort study. *Clin Psychol Sci.* 2016; Mar 22:2167702616630958.
[Google Scholar](#)
- 21.** Robins LN, Helzer JE, Croughan J, Ratcliff KS. National Institute of Mental Health Diagnostic Interview Schedule: its history, characteristics, and validity. *Arch Gen Psychiatry.* 1981;38(4):381-389.

[Article](#)

[Article](#) | [PubMed](#) | [Google Scholar](#) | [Crossref](#)

- 22.** Robins LN, Cottler L, Bucholz KK, Compton W. *Diagnostic Interview Schedule for DSM-IV.* St Louis, MO: Washington University School of Medicine; 1995

- icine, 1995.
- 23.** American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 3rd ed, revised. Washington, DC: American Psychiatric Association; 1987.
 - 24.** American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. ed 4. Washington, DC: American Psychiatric Association; 1994.
 - 25.** Israel S, Moffitt TE, Belsky DW, et al. Translating personality psychology to help personalize preventive medicine for young adult patients. *J Pers Soc Psychol*. 2014;106(3):484-498.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
 - 26.** Thomson WM, Poulton R, Broadbent JM, et al. Cannabis smoking and periodontal disease among young adults. *JAMA*. 2008;299(5):525-531.

[Article](#)

[Article](#) | [PubMed](#) | [Google Scholar](#) | [Crossref](#)

- 27.** Hancox RJ, Poulton R, Greene JM, et al. Systemic inflammation and lung function in young adults. *Thorax*. 2007;62(12):1064-1068.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 28.** Pletcher MJ, Vittinghoff E, Kalhan R, et al. Association between marijuana exposure and pulmonary function over 20 years. *JAMA*. 2012;307(2):173-181.

[Article](#)

[Article](#) | [PubMed](#) | [Google Scholar](#) | [Crossref](#)

- 29.** Hancox RJ, Poulton R, Ely M, et al. Effects of cannabis on lung function: a population-based cohort study. *Eur Respir J*. 2010;35(1):42-47.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)

- 30.** Costello EJ, Copeland WE, Shanahan L, Worthman CM, Angold A. C-reactive protein and substance use disorders in adolescence and early adulthood: a prospective analysis. *Drug Alcohol Depend.* 2013;133(2):712-717.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 31.** Vlassopoulos A, Lean ME, Combet E. Influence of smoking and diet on glycated haemoglobin and "pre-diabetes" categorisation: a cross-sectional analysis. *BMC Public Health.* 2013;13:1013.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 32.** Bergström J, Eliasson S, Dock J. A 10-year prospective study of tobacco smoking and periodontal health. *J Periodontol.* 2000;71(8):1338-1347.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 33.** Houston TK, Person SD, Pletcher MJ, Liu K, Iribarren C, Kiefe CI. Active and passive smoking and development of glucose intolerance among young adults in a prospective cohort: CARDIA study. *BMJ.* 2006;332(7549):1064-1069.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 34.** Mendall MA, Patel P, Ballam L, Strachan D, Northfield TC. C reactive protein and its relation to cardiovascular risk factors: a population based cross sectional study. *BMJ.* 1996;312(7038):1061-1065.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 35.** Craig WY, Palomaki GE, Haddow JE. Cigarette smoking and serum lipid and lipoprotein concentrations: an analysis of published data. *BMJ.* 1989;298(6676):784-788.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 36.** Zeng J, Williams SM, Fletcher DJ, et al. Reexamining the association between smoking and periodontitis in the Dunedin study with an enhanced analytical approach. *J Periodontol.* 2014;85(10):1390-1397

nanized analytical approach. *J Periodontol.* 2014;85(10):1350-1357.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)

37. Smit E, Crespo CJ. Dietary intake and nutritional status of US adult marijuana users: results from the Third National Health and Nutrition Examination Survey. *Public Health Nutr.* 2001;4(3):781-786.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

38. Vidot DC, Prado G, Hlaing WM, Arheart KL, Messiah SE. Emerging issues for our nation's health: the intersection of marijuana use and cardiometabolic disease risk. *J Addict Dis.* 2014;33(1):1-8.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

39. Ngueta G, Bélanger RE, Laouan-Sidi EA, Lucas M. Cannabis use in relation to obesity and insulin resistance in the Inuit population. *Obesity (Silver Spring).* 2015;23(2):290-295.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

40. Silvestri C, Di Marzo V. The endocannabinoid system in energy homeostasis and the etiopathology of metabolic disorders. *Cell Metab.* 2013;17(4):475-490.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

41. Després JP, Golay A, Sjöström L; Rimonabant in Obesity-Lipids Study Group. Effects of rimonabant on metabolic risk factors in overweight patients with dyslipidemia. *N Engl J Med.* 2005;353(20):2121-2134.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

42. Pi-Sunyer FX, Aronne LJ, Heshmati HM, Devin J, Rosenstock J; RIO-North America Study Group. Effect of rimonabant, a cannabinoid-1 receptor blocker, on weight and cardiometabolic risk factors in overweight or obese patients: RIO-North America: a randomized controlled trial [published correction appears in *JAMA*. 2006;295(11):1252]. *JAMA.* 2006;295(7):761-775.

- 43.** McPartland JM, Duncan M, Di Marzo V, Pertwee RG. Are cannabidiol and Δ^9 -tetrahydrocannabinol negative modulators of the endocannabinoid system? a systematic review. *Br J Pharmacol*. 2015;172(3):737-753.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 44.** Muniyappa R, Sable S, Ouwerkerk R, et al. Metabolic effects of chronic cannabis smoking. *Diabetes Care*. 2013;36(8):2415-2422.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 45.** Rodondi N, Pletcher MJ, Liu K, Hulley SB, Sidney S; Coronary Artery Risk Development in Young Adults (CARDIA) Study. Marijuana use, diet, body mass index, and cardiovascular risk factors (from the CARDIA study). *Am J Cardiol*. 2006;98(4):478-484.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 46.** Lee MH, Hancox RJ. Effects of smoking cannabis on lung function. *Expert Rev Respir Med*. 2011;5(4):537-546.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 47.** Mittleman MA, Lewis RA, Maclure M, Sherwood JB, Muller JE. Triggering myocardial infarction by marijuana. *Circulation*. 2001;103(23):2805-2809.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 48.** Mukamal KJ, Maclure M, Muller JE, Mittleman MA. An exploratory prospective study of marijuana use and mortality following acute myocardial infarction. *Am Heart J*. 2008;155(3):465-470.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 49.** Thomas G, Kloner RA, Rezkalla S. Adverse cardiovascular, cerebrovascular, and peripheral vascular effects of marijuana inhalation: what cardiolo-

gists need to know. *Am J Cardiol.* 2014;113(1):187-190.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

- 50.** Jouanjus E, Lapeyre-Mestre M, Micallef J; French Association of the Regional Abuse and Dependence Monitoring Centres (CEIP-A) Working Group on Cannabis Complications. Cannabis use: signal of increasing risk of serious cardiovascular disorders. *J Am Heart Assoc.* 2014;3(2):e000638. doi:[10.1161/JAHA.113.000638](#).
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 51.** Kinane DF, Chestnutt IG. Smoking and periodontal disease. *Crit Rev Oral Biol Med.* 2000;11(3):356-365.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 52.** Wolff V, Lauer V, Rouyer O, et al. Cannabis use, ischemic stroke, and multifocal intracranial vasoconstriction: a prospective study in 48 consecutive young patients. *Stroke.* 2011;42(6):1778-1780.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 53.** Ducros A. Reversible cerebral vasoconstriction syndrome. *Lancet Neurol.* 2012;11(10):906-917.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 54.** ElSohly MA, Mehmedic Z, Foster S, Gon C, Chandra S, Church JC. Changes in cannabis potency over the last two decades (1995-2014): analysis of current data in the United States. *Biol Psychiatry.* 2016;79(7):613-619.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 55.** Silins E, Horwood LJ, Patton GC, et al; Cannabis Cohorts Research Consortium. Young adult sequelae of adolescent cannabis use: an integrative analysis. *Lancet Psychiatry.* 2014;1(4):286-293.
[PubMed](#) | [Google Scholar](#) | [Crossref](#)
- 56.** Hancox PJ, Shin HH, Gray AB, Boulton P, Sears MP. Effects of quitting

56. Hancock KS, Shim TM, Gray AK, Poulton R, Sears MK. Effects of quitting cannabis on respiratory symptoms. *Eur Respir J*. 2015;46(1):80-87.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

57. Arseneault L, Cannon M, Poulton R, Murray R, Caspi A, Moffitt TE. Cannabis use in adolescence and risk for adult psychosis: longitudinal prospective study. *BMJ*. 2002;325(7374):1212-1213.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

58. Meier MH, Hill ML, Small PJ, Luthar SS. Associations of adolescent cannabis use with academic performance and mental health: a longitudinal study of upper middle class youth. *Drug Alcohol Depend*. 2015;156:207-212.

[PubMed](#) | [Google Scholar](#) | [Crossref](#)

✓ JAMA Psychiatry

✓ JAMA Network™

✓ Help



Get the latest from JAMA Psychiatry



[Privacy Policy](#) | [Terms of Use](#)

[Sign Up](#)



© 2020 American Medical Association. All Rights Reserved.

[Terms of Use](#) | [Privacy Policy](#) | [Accessibility Statement](#)

POWERED BY  SILVERCHAIR
INFORMATION/SYSTEMS