


Ribosomal Proteins Rpl22 and Rpl22I1 Control Morphogenesis by Regulating Pre-mRNA Splicing

Graphical Abstract



Authors

Yong Zhang, Monique N. O'Leary, Suraj Peri, ..., David R. Morris, Brian K. Kennedy, David L. Wiest

Correspondence

david.wiest@fccc.edu

In Brief

Zhang et al. reveal that ribosomal proteins can perform regulatory functions of fundamental importance while separated from the ribosome. They demonstrate that ribosomal proteins Rpl22 and Rpl22-Like1 (Like1) are retained in the nucleus during gastrulation, where they control morphogenesis by modulating the splicing of Smad2 pre-mRNA in cooperation with HNRNP-A1.

Highlights

- Ribosomal proteins Rpl22 and Like1 antagonistically control morphogenesis
- During gastrulation, Rpl22 and Like1 act away from the ribosome in the nucleus
- Rpl22 and Like1 control gastrulation by regulating the splicing of Smad2 pre-mRNA
- Control of gastrulation by Rpl22 and Like1 involves cooperation with HNRNP-A1



Ribosomal Proteins Rpl22 and Rpl22l1 Control Morphogenesis by Regulating Pre-mRNA Splicing

Yong Zhang,^{1,6} Monique N. O'Leary,^{2,6} Suraj Peri,¹ Minshi Wang,¹ Jikun Zha,¹ Simon Melov,² Dietmar J. Kappes,¹ Qing Feng,³ Jennifer Rhodes,¹ Paul S. Amieux,⁴ David R. Morris,⁵ Brian K. Kennedy,² and David L. Wiest^{1,7,*}

¹Blood Cell Development and Function Program, Fox Chase Cancer Center, Philadelphia, PA 19111, USA

²Buck Institute for Research on Aging, Novato, CA 94945, USA

³Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA

⁴Department of Pharmacology

⁵Department of Biochemistry

University of Washington, Seattle, WA 98195, USA

⁶Co-first author

⁷Lead Contact

*Correspondence: david.wiest@fccc.edu

<http://dx.doi.org/10.1016/j.celrep.2016.12.034>

SUMMARY

Most ribosomal proteins (RP) are regarded as essential, static components that contribute only to ribosome biogenesis and protein synthesis. However, emerging evidence suggests that RNA-binding RP are dynamic and can influence cellular processes by performing “extraribosomal,” regulatory functions involving binding to select critical target mRNAs. We report here that the RP, Rpl22, and its highly homologous paralog Rpl22-Like1 (Rpl22l1 or Like1) play critical, extraribosomal roles in embryogenesis. Indeed, they antagonistically control morphogenesis through developmentally regulated localization to the nucleus, where they modulate splicing of the pre-mRNA encoding *smad2*, an essential transcriptional effector of Nodal/TGF- β signaling. During gastrulation, Rpl22 binds to intronic sequences of *smad2* pre-mRNA and induces exon 9 skipping in cooperation with hnRNP-A1. This action is opposed by its paralog, Like1, which promotes exon 9 inclusion in the mature transcript. The nuclear roles of these RP in controlling morphogenesis represent a fundamentally different and paradigm-shifting mode of action for RP.

INTRODUCTION

Ribosomal proteins (RP) are basic components of the ribosome that are generally thought to contribute to the assembly of the ribosome and its ability to synthesize proteins. Inactivation of RP has been linked to clinical syndromes collectively known as “ribosomopathies” (Narla and Ebert, 2010). Ribosomopathies exhibit not only impaired erythropoiesis and increased risk for development of leukemia but also other abnormalities in a variety of organ systems (Narla and Ebert, 2010). These developmental

anomalies include short stature, craniofacial defects, thumb malformation, urogenital abnormalities, and heart defects, which strongly suggests that RP might play a role in embryogenesis (Narla and Ebert, 2010). A recent example of this is the finding that haploinsufficiency of *RPSA* causes human asplenia, suggesting that RP inactivation can cause distinct and tissue-restricted developmental abnormalities (Bolze et al., 2013). Mutations of RP have been shown in many cases to impair ribosome biogenesis, which activates p53 by attenuating the function of the p53 ubiquitin ligase MDM2 (Zhang and Lu, 2009). However, emerging evidence has begun to reveal that certain RP may also perform specialized regulatory roles in biological processes through p53-independent “extraribosomal functions,” acting to regulate context-dependent translation or transcription, and doing so from outside of the ribosome (Warner and McIntosh, 2009; Xue and Barna, 2012). For example, extraribosomal Rpl26 and Rpl13a can modulate the translation of selected target RNAs by binding to their 5' and 3' UTRs, respectively (Mukhopadhyay et al., 2008; Takagi et al., 2005). Likewise, Rps3 and Rps14 have been shown to interact with DNA-binding complexes and regulate gene-specific transcription (Wan et al., 2007; Zhou et al., 2013).

We previously established that neither of the highly homologous RP paralogs Rpl22 or Rpl22-Like1 (Rpl22l1 or Like1) is required for general protein synthesis; however, these paralogs do play distinct, antagonistic, regulatory roles in blood cell development (Anderson et al., 2007; Rao et al., 2012; Zhang et al., 2013). Indeed, the ability of Like1 to promote the emergence of embryonic hematopoietic stem cells (HSC) is dependent upon its ability to directly bind and facilitate the translation of mRNA encoding the essential transcription factor, *smad1*. In contrast, Rpl22 acts in direct opposition to repress *smad1* translation (Zhang et al., 2013). These findings revealed that the RNA-binding RP, Rpl22 and Like1, play an important regulatory role in hematopoiesis, from outside of the ribosome (Zhang et al., 2013). Our previous analysis in zebrafish also revealed that Rpl22 and Like1 are abundantly expressed throughout early development from the two-cell stage to 18 hr post-fertilization (hpf) (Zhang et al., 2013), raising the possibility that they might also act to

regulate early development, as has been found for other molecular effectors exhibiting similar expression patterns (Langdon and Mullins, 2011; Schier, 2007).

Here, we used antisense morpholinos (MO) targeting the ATG translational start codons of Rpl22 and Like1 to attenuate the translation of both maternal and zygotic mRNA encoding these proteins to repress their expression and assess the impact on development of zebrafish embryos. We showed that Like1 knockdown disrupted the convergence and extension (C&E) phase during gastrulation and, consequently, elaboration of the normal body plan. The molecular basis for disruption of C&E is that splicing of pre-mRNA encoding *smad2*, an essential mediator of Nodal/TGF- β signaling, was impaired, revealing nuclear functions for Rpl22 and Like1 in regulating pre-mRNA splicing. Indeed, Like1 knockdown resulted in skipping of exon 9 of *smad2*, which blocked protein expression. The skipping of exon 9 was caused by Rpl22, as both the mis-splicing of *smad2* and the defect in C&E observed in Like1 morphants were rescued by simultaneously knocking down Rpl22. The ability of Rpl22 and Like1 to control *smad2* pre-mRNA splicing coincided with a developmentally controlled retention of these proteins in the nucleus and was associated with direct binding of Rpl22 to a consensus motif in *smad2* intron 8, immediately preceding skipped exon 9. Using RNA sequencing (RNA-seq) we determined that numerous other pre-mRNAs were mis-spliced in Like1 morphants and these targets shared the features of having consensus Rpl22/Like1 binding sites in the intron preceding the skipped exon and potential binding sites for splicing modulator hnRNP-A1. hnRNP-A1 function is required for the ability of Rpl22 to disrupt *smad2* splicing, as exon 9 inclusion is restored upon knockdown of hnRNP-A1 in Like1 morphants. Together, these data reveal that the RP Rpl22 and Like1 antagonistically control gastrulation through a nuclear role in regulating mRNA splicing, in zebrafish as well as in mammals.

RESULTS

Rpl22 and Like1 Play Distinct and Antagonistic Roles during Gastrulation

We previously demonstrated that neither of the highly homologous RP paralogs, Rpl22 and Like1, is required for general protein synthesis; however, these paralogs do play distinct, antagonistic, regulatory roles in hematopoiesis that are mediated by binding to and controlling the translation of *smad1* mRNA (Anderson et al., 2007; Rao et al., 2012; Shen et al., 2012; Zhang et al., 2013). Moreover, we showed that the mRNAs encoding Rpl22 and Like1 were abundantly expressed in zebrafish embryos beginning at the two-cell stage (Zhang et al., 2013). To determine if Rpl22 and Like1 regulate morphogenesis, we used antisense MO that target the ATG start codons and repress the translation of both maternal and newly synthesized zygotic *rpl22* and *rpl22l1* (Like1) mRNA (Zhang et al., 2013) (Figures S1A and S1B), enabling analysis of the role of these proteins in early development. Like1 morpholino-treated embryos (Like1 morphants) exhibited defects in anterior-posterior extension during late gastrulation (Figure 1A), as indicated by the reduced angle between the anterior and posterior ends (Figure 1B). Importantly, whereas knockdown of Rpl22 did not disrupt exten-

sion (Figures 1A and 1B), simultaneous knockdown of Rpl22 and Like1 (double morphants [D-MOs]) completely corrected the extension defect observed upon knockdown of Like1 alone (Figures 1A, right, and 1B). This indicates that Like1 plays a critical role in promoting C&E during gastrulation and that its ability to promote C&E is antagonized by its paralog, Rpl22. Co-injecting Like-MO and mCherry-CAAX (membrane-targeted mCherry) mRNA revealed that knockdown of Like1 caused a broadening of the notochord in 10 hpf Like1 morphants (Figure 1C), indicating that the convergence of notochord precursors was also impaired by Like1 knockdown. The shortening and widening of mesoderm (*ntl*, *gsc*), endoderm (*sox32/sox17*), paraxial mesoderm (*papc*), adaxial (*myod1*), and brain and neural tissue (*six3*, *pax2*, *krox20*, and *hgg1/dlx3b*) in Like1 morphants was confirmed by whole-mount in situ hybridization (WISH) using probes for markers of those tissues (Figures 1D, 1E, and S1C–S1J). To gain insight into the molecular basis for the disruption of C&E during gastrulation, we assessed whether signaling pathways previously determined to be critical for early morphogenesis were disrupted. We found that signaling through the Bmp/pSmad5, canonical and noncanonical Wnt, PI3-Kinase/Akt, and Stat3/Liv1/e-cadherin pathways controlling migration of developing tissues was unaffected by Like1 knockdown (Figures S1K–S1P) (Heisenberg and Solnica-Krezel, 2008; Kimelman and Griffin, 2000; Solnica-Krezel, 2005). However, the expression of *lefty1* was essentially eliminated in Like1 morphants at 16 hpf (Figure 1F). Because *lefty1* is a direct target of Nodal/Smad2 signaling, we assessed whether Smad2 signaling was altered in Like1 morphants (Smith et al., 2011; Stemple, 2000; Thisse et al., 2000). Indeed, both Smad2 phosphorylation and expression were reduced between 10 and 16 hpf (Figures 1G and 1H). Smad2 signaling plays an essential role in regulating gastrulation and later morphogenesis (Heyer et al., 1999; Nomura and Li, 1998; Waldrip et al., 1998). Interestingly, maternal Smad2/Nodal signaling and expression of the downstream target *squint* (*sqt*) were intact at 4.7 hpf, prior to the onset of gastrulation (Figures 1I and S1Q), indicating that the ability of Like1 to control Smad2 expression was developmentally regulated.

Smad2 Splicing Is Regulated by Rpl22 and Like1 during Gastrulation

To determine how Like1 knockdown repressed the expression of zygotic Smad2, we tested whether *smad2* mRNA levels were altered. Interestingly, while total *smad2* mRNA levels were unchanged, most of the *smad2* mRNA was significantly smaller in size in 10 hpf Like1 morphants (Figure 2A). The alteration of *smad2* mRNA size was not observed until 6 hpf, after the switch from maternal to zygotic transcription (Kane and Kimmel, 1993) (Figure S2A). Sequence analysis revealed that the basis for the reduction in *smad2* mRNA size was that it lacked exon 9, such that exon 8 was directly spliced to exon 10 (Figures 2B and 2C). Although knockdown of Rpl22 alone did not affect *smad2* splicing, simultaneous knockdown of Rpl22 with Like1 significantly reduced the *smad2* mis-splicing observed in Like1 morphants, suggesting antagonistic regulation by these RP paralogs (Figure 2A, right lane). Isoform-specific qPCR confirmed that normal *smad2* splicing was increased from ~35% in Like1 morphants to ~70% in morphants where Rpl22 was knocked




Figure 1. Opposing Roles of RP Paralogs Rpl22 and Like1 in Regulating Gastrulation

(A and B) One-cell stage zebrafish embryos were injected with translational-blocking Like1-A-MO (L1-A-MO; 2 ng), L22-A-MO (6 ng), or both (D-MOs), following which effects on gastrulation were assessed. The images of embryos represent lateral views at 10 hpf. The red and black lines indicate the anterior and posterior ends of the body axis, respectively. The angle (θ), which defines the degree of extension, was measured between the red arrow and dashed black line and is represented graphically as the mean \pm SD. Control (Cont.; black); Like1 MO (L1-A-MO; red); Rpl22 MO (L22-A-MO; yellow); and double-morphants (D-MO; green). Triplicate samples were quantified and the mean \pm SD is depicted graphically. ** $p < 0.01$.

(C) Imaging of notochord in Like1 morphants (10 hpf) co-injected with 100 pg mCherry-CAAX mRNA at the one-cell stage. The dorsal view, anterior is at the top. The lateral notochord boundaries are indicated by the dotted yellow lines, and the width of notochord was marked by white lines.

(D and D') Expression patterns of *ntl* in 10 hpf Like1 morphants. Red arrows mark changes in distribution in the images representing lateral (D) and dorsal (D') views.

(E and F) *sox32* and *lefty1* expression in Like1 morphants. Red arrows indicate changes in expression or distribution. (E) 75%-epiboly stage, dorsal view. (F) Lateral view, 16 hpf.

(G–I) Phospho-Smad2 and total-Smad2 were assessed in 10 hpf (G), 16 hpf (H), and 4.7 hpf (I) Like1 morphants by immunoblotting.

All results are representative of at least three experiments performed. See also Figure S1.

down together with Like1 (i.e., in double-morphants [D-MOs]; Figure 2D).

The C&E Defects in Like1 Morphants Result from Smad2 Mis-Splicing

To assess whether *smad2* pre-mRNA mis-splicing was responsible for the C&E defects, we sought to specifically replicate the mis-splicing of *smad2* using MO targeting the *smad2* intron 8-exon 9 boundary (S2-i8e9-MO; Figure 3A). We verified that the S2-i8e9-MO replicates the mis-splicing of *smad2* and reduces Smad2 protein levels (Figures 3B, S2B, and S2C). Moreover, the S2-i8e9-MO also phenocopied the morphological defects and impaired *lefty1* expression observed in Like1 morphants (Figures 3C–3E). The mis-spliced *smad2* transcript appeared to be disrupting development by reducing Smad2 expression rather than functioning as a dominant negative. Indeed, ectopic expression of the mis-spliced *smad2* transcript failed to block Smad2-dependent signaling events induced by a constitutively activated Alk4 mutant (*Tar**) (Figures S3A and S3B). This is not unexpected because, despite the retention of the translational reading frame, we failed to detect a truncated protein produced from the exon 9-skipped *smad2* mRNA, either

endogenously or upon ectopic expression of the mutant cRNA (Figures S3C and S3D). The inability to produce a stable, truncated protein has also been reported for a murine *Smad2* splice variant lacking exons 9 and 10, in which the translational reading frame was preserved (Liu et al., 2004). Having found that the mis-splicing of *smad2* was sufficient to phenocopy the C&E defects observed in Like1 morphants, we next asked if restoring Smad2 signaling could rescue those defects. Indeed, ectopic expression of constitutively active Smad2 (Ca-Smad2) did alleviate the C&E defects in Like1 morphants (morphology and *ntl*/*myod1* distribution; Figures 3F–3H) (Dick et al., 2000). Taken together, these results suggest that Rpl22 and Like1 antagonistically regulate C&E by controlling the splicing of *smad2*, a critical regulator of gastrulation.

The Subcellular Localization of Rpl22 and Like1 in Zebrafish Embryos Is Developmentally Regulated

For Rpl22 and Like1 to directly regulate *smad2* pre-mRNA splicing during gastrulation, Rpl22 and Like1 would have to be located in the nucleus. Indeed, immunofluorescence analysis of ectopically expressed HA-tagged RP revealed that Like1 and Rpl22 were located primarily in the nucleus at 6 and 10




Figure 2. The Gastrulation Defects in Like1 Morphants Result from Skipping of *smad2* Exon 9

(A and B) RT-PCR analysis of *smad2* mRNA in Like1 (L1-A-MO), Rpl22 (L22-A-MO), and double morphants (D-MOs). *smad2* mRNA was evaluated by RT-PCR using primers (black arrowheads) amplifying the sequences between exons 5–10.

(C) Like1 knockdown causes skipping of *smad2* exon 9. Sequence analysis of the smaller *smad2* mRNA species caused by Like1 knockdown (red arrow in A).

(D) Quantitative RT-PCR analysis of the relative expression of intact *smad2* mRNA (blue arrow, left). The blue and red lines identify the positions of real-time primers used to detect intact *smad2* mRNA. Triplicate samples were quantified, and the mean \pm SD is depicted graphically. P values are indicated. All results are representative of at least three experiments performed. See also Figure S2.

hpf (Figures 4A and S4A). The nuclear localization of Like1 and Rpl22 appears to be developmentally regulated, as they relocalized to the cytoplasm at 24 hpf, after gastrulation and morphogenesis are complete (Figure 4B). Importantly, at 24 hpf, Like1 knockdown no longer caused *smad2* mis-splicing (Figure 4C). The nuclear retention of Rpl22 and Like1 requires both their NLS motifs and their ability to bind RNA (Figures S4B–S4E). These results indicate that Rpl22 and Like1 regulate *smad2*

pre-mRNA splicing in a developmentally controlled manner, and their ability to do so is tightly linked to nuclear localization during gastrulation.

Like1 Deficiency Disrupts *Smad2* Splicing in Mouse Embryos

To determine if these RP also regulate *Smad2* splicing during mammalian gastrulation, we generated mice in which the *Rpl22l1* gene was ablated (Figures 5A–5C). Intercrossing *Rpl22l1*^{+/-} mice revealed that Like1 deficiency was embryonic lethal, as no *Rpl22l1*^{-/-} mice were observed among 54 offspring. Moreover, *Rpl22l1*^{-/-} embryos were absent embryonically by 12 days post-coitus (dpc) and displayed aberrant morphology at 9.5 dpc (Figure 5D). When the status of *Smad2* pre-mRNA splicing was assessed at 6.5 dpc, we observed mis-splicing of the *Smad2* pre-mRNA (Figures 5E and 5F); however, unlike the mis-splicing observed in zebrafish, both exons 7 and 8 of murine *Smad2* mRNA were skipped in the *Rpl22l1*^{-/-} embryos (Figure 5G). Together, these data indicate that the regulation of *Smad2* splicing during gastrulation is not only observed in zebrafish but is also conserved in mammals.

Like1 Knockdown Disrupts the Splicing of Numerous Pre-mRNA Targets

To gain insight into how Rpl22 and Like1 might antagonistically regulate *smad2* splicing, we performed RNA-seq to identify the set of pre-mRNAs whose splicing they regulate. RNA-seq analysis of Like1 morphants revealed more than 300 mis-spliced targets, including *smad2* (Figure 6A). Pathway analysis revealed that the mis-spliced targets were enriched for those involved in DNA replication, morphogenesis, and regulation of BMP signaling (Figure 6B). Most of the mis-splicing events represented exon skipping, with a few instances of alternative exon usage (Figure 6C). Moreover, MaxEntScan analysis of the mis-spliced targets revealed that the skipped exons (Figure 6D; skipped [S]) exhibited weaker 5' splice donor and 3' splice acceptor sites than the included exons in the same genes (Figure 6D; not skipped [NS]) (Lu et al., 2013). Hence, Like1 knockdown preferentially induced the skipping of exons with weak splice sites, suggesting that additional splicing factors were necessary for recognition of these exons by the spliceosome (Lopez, 1998). It is well established that *trans*-acting splicing factors such as hnRNP A/B family members are able to preferentially bind to exonic splicing silencers (ESS) and antagonize the binding of SR proteins to exonic splicing enhancers (ESE), thereby leading to exon skipping (Wang and Burge, 2008; Zhu et al., 2001). The observation that most of the mis-splicing events in Like1 morphants were exon skips raised the possibility that the exonic elements involved might be bound by *trans*-acting factors that could influence their function and cause exon skipping (Wang et al., 2004). To test this hypothesis, we validated a set of mis-spliced target genes by RT-PCR (Figure S5) and interrogated the sequences surrounding the skipped exons to identify common features. The FAS-ESS and RESCUE-ESE algorithms were used to predict potential ESS or ESE (Wang et al., 2004). Indeed, we discovered that G-rich (GGGG or GGG) motifs were enriched in skipped exons (Figure S6A). Furthermore, the consensus Rpl22/like1 stem-loop binding motif was found in




Figure 3. The C&E Defects in Like1 Morphants Can Be Rescued by Re-establishing *smad2* Signaling

(A) Schematic of the morpholino used to induce exon 9 skipping (S2-i8e9-MO).

(B) Immunoblotting of detergent extracts reveals a reduction in total and phospho-Smad2 protein expression in the S2-i8e9-morphants.

(C–E) S2-i8e9-MO induction of *smad2* mis-splicing phenocopied the C&E defects caused by Like1 knockdown, as indicated by altered morphology (C, red arrow) and alterations in *ntl*/*hgg1*/*dlx3b* and *lefty1* expression and distribution, as measured by in situ hybridization (D, 10 hpf, red arrows, anterior dorsal view; E, 16 hpf, lateral view).

(F–H) mRNA encoding constitutively activated *smad2* (Ca-Smad2, 20 pg) was used for injection alone or co-injected with Like1-A-MO. Embryo morphology (F) as well as the abnormal distribution of *ntl* (G, red arrows) and *myoD* (H, red arrows) at 10 hpf can be rescued by ectopic expression of Ca-Smad2. All embryos are dorsal view with the anterior on top at 10 hpf.

All results are representative of at least three experiments performed. See also Figure S3.

introns immediately preceding the skipped exons (Figure 6E), suggesting that Rpl22/Like1 can directly bind to target pre-mRNAs and regulate their splicing. Importantly, these features were also found in *smad2* pre-mRNA (Figure S6B). Thus, these data support a model where Rpl22 and Like1 antagonistically regulate gastrulation by directly binding to pre-mRNA targets, including *smad2*, and promoting their mis-splicing in conjunction with a *trans*-acting factor(s) that recognizes a G-rich motif (Figure 6F).

The Ability of Rpl22 to Disrupt *smad2* Pre-mRNA Splicing in Like1 Morphants Is Dependent upon hnRNP-A1

To test this model, we used in vivo RNA:protein cross-linking immunoprecipitation (CLIP) analysis to determine if Rpl22 and Like1 can bind their consensus motif in *smad2* pre-mRNA. Both Rpl22 and Like1, but not their m88 RNA-binding mutant forms, were able to bind to intron 8 of *smad2* pre-mRNA, which immediately precedes the skipped exon 9 (Figures 7A and 7B), suggesting that the regulation of *smad2* splicing by Rpl22 and Like1 is direct. We next sought to identify the *trans*-acting factor(s) that bind the G-rich motif present in the pre-mRNAs mis-spliced in Like1 morphants. hnRNP-A1 (A1) has been reported to recognize a motif similar to the G-rich sequence motif observed in the targets mis-spliced in Like1 morphants, and zebrafish A1 is expressed and localized in the nucleus during gastrulation (Figure S7A) (Gabut et al., 2008). Moreover, A1 is a well-established splicing modulator that can bind ESS and inhibit exon inclusion (Zhu et al., 2001). To determine if A1 was contributing to the ability of Rpl22 and Like1 to modulate *smad2* splicing, we overexpressed A1 mRNA and found that A1, but not other hnRNP, such as hnRNP-H and hnRNP-I (Y.Z., unpublished data), induced *smad2* exon 9 skipping (Figure 7C, red arrow). Moreover, knocking down A1 (validated in Figure S7B) in

Like1 morphants significantly reduced the skipping of *smad2* exon 9 in Like1 morphants (Figure 7D, red arrow). Finally, knock-down of both Rpl22 and A1 completely suppressed the *smad2* mis-splicing normally observed in Like1 morphants (Figure 7D, red arrow), suggesting that Rpl22 and A1 collaborate in promoting *smad2* mis-splicing in the absence of Like1. hnRNPs have been reported to physically and specifically interact with selected RPs (Kristensen et al., 2012). To determine if the ability of Rpl22 and A1 to promote *smad2* mis-splicing involved their physical association, we performed co-precipitation analysis. We found that A1 co-precipitated with Rpl22, but not Like1, in detergent extracts of 10 hpf zebrafish embryos, demonstrating a physical interaction between Rpl22 and A1 (Figure 7E). Importantly, the Rpl22-A1 association was not mediated by an RNA bridge, as it was not disrupted by RNase treatment (Figure 7E).

DISCUSSION

Mutations in RP have long been linked to developmental anomalies manifested among the group of inherited syndromes collectively known as ribosomopathies. Although the notion that these anomalies result from generalized perturbations of ribosome biogenesis or function is increasingly viewed as too simplistic, little insight has been gained into the molecular basis for their genesis. We report here nuclear functions for Rpl22 and Like1, which when disrupted perturb embryonic patterning by impairing C&E during gastrulation. Indeed, Rpl22 and Like1 antagonistically control C&E through their developmentally regulated retention in the nucleus and their ability to influence the splicing of pre-mRNAs of key regulators of gastrulation, principally Smad2. The outcome of these splicing events is determined by the antagonistic balance of Rpl22 and Like1 and entails cooperation with the splicing modulator hnRNP-A1. These findings reveal a fundamentally different way to view the function of




Figure 4. The Regulation of *smad2* Pre-mRNA Splicing during Gastrulation by Rpl22 and Like1 Is Associated with Their Retention in the Nucleus

(A and B) Subcellular location of epitope-tagged Rpl22 and Like1 at 10 hpf (A) and 24 hpf (B). mRNA (100 pg) encoding HA-zRpl22 (HA/L22) and HA-zLike1 (HA/L1) was co-injected with mRNA encoding mCherry-CAAX into one-cell stage embryos and visualized with HA antibody immunostaining. mCherry-CAAX marked the cell membrane, and DAPI marked the nucleus. The red scale bar represents 10 μ m.

(C) RT-PCR detection of *smad2* mis-splicing. Following Like1 MO injection, *smad2* mis-splicing was assessed by RT-PCR in 10 and 24 hpf Like1 morphants.

All results are representative of at least three experiments performed. See also Figure S4.

RP, in that they are capable of not only existing independent of intact ribosomes but are capable, in that form, of exerting a profound influence on critical events during development in zebrafish as well as in mammals. These findings also raise a number of crucial questions relating to how Rpl22 and Like1 are retained in the nucleus, the molecular basis by which they regulate the splicing of pre-mRNA targets, and how Rpl22 and Like1 are able to exert opposing effects on splicing.

Because the control of splicing by Rpl22 and Like1 is linked to their retention in the nuclear, the effects on splicing are clearly

mediated in an extraribosomal fashion; however, the basis for the developmentally controlled retention of Rpl22 and Like1 in the nucleus remains a critical unanswered question. There are two likely explanations. First, Rpl22 and Like1 may assemble into the ribosome in the nucleolus, traffic to the cytosol, and then separate from the ribosome in the cytosol, following which they traffic back to the nucleus. The separation of Rpl22 and Like1 from the ribosome could be induced by post-translational modifications. This has been observed for Rpl13a, which is displaced from the ribosome by interferon-mediated phosphorylation. Upon release, Rpl13a is able to bind to cytosolic mRNA species and silence their translation as part of the GAIT complex (Mukhopadhyay et al., 2008). Motif prediction analysis suggests that both Rpl22 and Like1 possess numerous consensus phosphorylation sites; however, it remains unclear if any are actually used in vivo. Rpl22 has been reported to be modified by SUMO in *Drosophila*, which is required to localize Rpl22e to the nucleoplasm (Kearse et al., 2013). Nevertheless, because *Drosophila* Rpl22 contains a large, unique N-terminal extension, it remains unclear whether the SUMOylation of Rpl22 or Like1 plays a role in influencing their subcellular localization in other species. Rpl22 and Like1 might also be induced to dissociate from the ribosome through protein:protein interactions. We have shown that Rpl22 associates hnRNP-A1, a protein whose localization is reported to change dynamically during embryonic development (Vautier et al., 2001). Specifically, when zygotic transcription is activated, hnRNP-A1 accumulates in the nucleus (Vautier et al., 2001). The developmental changes in hnRNP-A1 localization, along with its genetic and physical interaction with Rpl22, raise the possibility that hnRNP-A1 may play a role in retention of Rpl22 in the nucleus during gastrulation. Nevertheless, hnRNP-A1 does not display the same association with Like1, suggesting that Like1 retention would be mediated by association with a different factor(s). Another mode of controlling the localization of Rpl22 and Like1 might entail direct trafficking to the nucleus after synthesis in the cytosol, through a process that does not involve their assembly onto the ribosome. We have shown that Rpl22 and Like1 localization in the nucleus requires both their NLS and RNA-binding motifs. Because the retention of Rpl22 and Like1 in the nucleus coincides with the switch to zygotic transcription during zebrafish embryogenesis, Rpl22 and Like1 retention in the nucleus is likely to be due, at least in part, to association with the large number of nascent transcripts being made in the nucleus during gastrulation. This is consistent with a previous report indicating that RP can associate with nascent RNAs in budding yeast (Schroder and Moore, 2005). Given the intimate connection between transcription and splicing, this represents a plausible explanation for developmental control of the retention of Rpl22 and Like1 in the nucleus and their involvement in regulating splicing.

The critical pre-mRNA target through which Rpl22 and Like1 modulation of splicing controls gastrulation is Smad2, a transcription factor whose function is essential for this process. Indeed, Like1 knockdown induces the skipping of *smad2* exon 9, which preserves the translational reading frame of the splice variant but nevertheless results in the loss of Smad2 protein. This results either from repression of translation, or more likely, from instability of the truncated protein product. The inability to




Figure 5. Effect of Like1 Deficiency on Smad2 Splicing during Murine Gastrulation

(A) Molecular strategy for targeted deletion of *Rpl22l1*. A Loxp-FRT-neo resistance-FRT cassette was inserted 3' to the first exon of *Rpl22l1*, and a second LoxP site was inserted 3' to the third exon. F1 heterozygous offspring were bred to *Mox2-cre* mice to delete the second and third exons of *Rpl22l1*, disrupting expression at the genomic locus.

(B and C) Strategy to genotype *Rpl22l1*^{+/L} or *Rpl22l1*^{+/-} mice. To genotype mice with LoxP sites flanking exons 2 and 3 of *Rpl22l1*, primers C and D were used to amplify a 223 or 263 bp product for wild-type (WT; *Rpl22l1*^{+/+}) and *Rpl22l1*-LoxP (*Rpl22l1*^{L/L}), respectively. Deletion of *Rpl22l1* is genotyped with primers B and D, which amplify a 607 bp product for the WT allele (*Rpl22l1*^{+/+}) and A and D, which amplify a 420 bp product for the mutant allele after cre recombination (*Rpl22l1*^{-/-}).

(D) Representative *Rpl22l1*^{-/-} embryo compared with *Rpl22l1*^{+/+} littermate control at 9.5 dpc.

(E–G) Effect of Like1 deficiency on splicing of *Smad2* pre-mRNA during murine gastrulation. Embryos derived from timed matings of *Rpl22l1*^{+/-} mice were isolated at 6.5 dpc (mid gastrulation), genotyped as above, and analyzed by RT-PCR using the indicated primers to identify alterations in *Smad2* splicing (E and F). Sequencing of the mis-spliced *Smad2* species found in *Rpl22l1*^{-/-} embryos revealed that it represented a species in which exon 6 was fused directly to exon 9, eliminating exons 7 and 8 (G).

All results are representative of at least three experiments performed.

generate a stable protein product appears to be linked to the particular exonic sequences lost (i.e., exon 9), because a *Smad2* variant lacking exon 3 is capable of supporting production of a truncated *Smad2* protein product (Dunn et al., 2005). Accordingly, the protein domain encoded by exon 9 is likely to play a critical role in maintaining the structural integrity of *Smad2*. Interestingly, while restoration of *Smad2* expression is able to alleviate the C&E defect observed in Like1 morphants, restoration of *Smad2* expression is not sufficient to rescue development beyond gastrulation, between 16 hpf and 3 dpf (data not shown). This is likely to be because the other targets that are mis-spliced upon Like1 knockdown are playing a role at more distal stages of development. The role of these mis-

spliced targets in later morphogenesis processes, such as heart development and left-right patterning, is currently under investigation.

Although it is clear that Rpl22 and Like1 regulate the splicing of numerous pre-mRNA targets in addition to *smad2*, it remains unclear how they do so. Rpl22 and Like1 could regulate splicing by directly binding to their pre-mRNA targets. Consistent with this possibility, we found that Rpl22 and Like1 can bind *smad2* pre-mRNA. Moreover, there are consensus Rpl22/Like1 binding motifs in the intron immediately preceding the skipped exon in all of validated, mis-spliced targets. Collectively, these data strongly suggest that Rpl22 and Like1, both RNA-binding proteins, are regulating splicing through direct binding to




Figure 6. Common Features of mRNA Targets Mis-spliced in Like1 Morphants

(A) *smad2* exon 9 skipping detected by RNA-seq. Alignment of RNA-seq reads to the genome reveals exclusion of *smad2* exon 9 in 10 hpf Like1 morphants (blue box).

(B) Gene Ontology analysis of transcripts affected by Like1 knockdown from RNA-seq analysis. Significant Gene Ontology terms ($p < 0.05$) are depicted as a bar graph with the p values represented as $-\log_{10}(p \text{ value})$ on the x axis.

(C) Schematic illustrating the type of alternative splicing identified by RNA-seq in 10 hpf Like1 morphants.

(D) Calculation of the strength of 5' and 3' splice sites of skipped (S) versus non-skipped (NS) exons in pre-mRNAs targeted in Like1 morphants. Results are represented as box and whisker plots. Statistical significance of the differences was determined using the Wilcoxon RS test.

(E) Schematic of the consensus hairpin bound by Rpl22/Like1.

(F) Schematic model of the common features of pre-mRNA targets affected by Like1 knockdown. Targets contained both a consensus Rpl22/Like1 binding motif in the preceding intron and a G-rich motif in the skipped exon.

RNA-seq analysis was performed on at least three independent biological replicates per condition. See also Figures S5 and S6.

pre-mRNA targets (Zhang et al., 2013). Alternatively, it should be noted that the zebrafish and human U2-snRNA, on which the U2-snRNP is assembled, also contains a consensus Rpl22/Like1 binding motif. Accordingly, it is also possible that Rpl22 and Like1 could influence splicing indirectly, through effects on the U2-snRNP, which plays a central role in RNA-splicing (Matera and Wang, 2014). Nevertheless, even if altered U2-snRNP function were contributing to mis-splicing, because consensus Rpl22 and Like1 binding sites were found in all validated targets, it re-

mains likely that direct binding of Rpl22 and Like1 to targets plays an important role.

One of the most interesting aspects of the regulation of development by the paralogs, Rpl22 and Like1, is the basis for their antagonistic functions. It was formerly thought that RP paralogs in lower organisms served largely redundant roles, but emerging evidence hints at functional specialization (Komili et al., 2007). Consistent with this notion, a few RP paralogs have been conserved in vertebrates, in which they exhibit some tissue




Figure 7. Factors through which Rpl22 and Like1 Regulate *smad2* Alternative Splicing

(A) Position of the real-time primers (red arrows) flanking the consensus Rpl22/Like1 binding site in intron 8 of *smad2* pre-mRNA.

(B) RNA-CLIP analysis of Rpl22/Like1 binding to *smad2* pre-mRNA. Embryos injected with mRNA encoding HA tagged Rpl22, Like1, or their RNA-binding mutants (m88) were harvested at 10 hpf. After light crosslinking, detergent nuclear extracts were immunoprecipitated using anti-HA antibody, and the co-precipitated RNA quantified by RT-PCR. Triplicate measurements are depicted graphically as mean \pm SD, and p values are indicated.

(C) Overexpression of hnRNP-A1 (A1) mRNA induces *smad2* mis-splicing. Embryos were injected with differing amounts of A1 mRNA, following which the effect on *smad2* mis-splicing was determined by RT-PCR at 10 hpf. The relative ratio of exon 9 skipped mRNA to intact *smad2* mRNA was quantified in triplicate and depicted graphically as the mean \pm SD.

(D) Genetic interaction of Rpl22 and A1. The ratio of exon 9 skipped to intact *smad2* was quantified in Like1 morphants in which A1 and/or Rpl22 was knocked down. The mean \pm SD of triplicate measurements was depicted graphically as in (C).

(E) Physical association between Rpl22, Like1 and A1. Anti-HA immunoprecipitation (IP) and anti-Flag immunoblots (IB) were performed on detergent extracts of embryos injected with 100 pg of mRNA encoding Flag-A1 and either HA-Rpl22 or HA-Like1, either before or after treatment with RNase.

All results are representative of at least three experiments performed. See also Figure S7.

restriction in their expression patterns, but their functions remain largely unexplored. We have previously reported that the antagonistic functions of Rpl22 and Like1 regulate the emergence of embryonic hematopoietic stem cells (Zhang et al., 2013). We now find that Rpl22/Like1 antagonism extends to the control of pre-mRNA splicing during gastrulation. Because Rpl22 and Like1 are more than 70% identical at the amino acid level, this raises the question of how proteins as highly homologous as these are able to perform antagonistic functions. The RNA-binding cores of Rpl22 and Like1 are nearly identical, and the helices that contact RNA are entirely identical and conserved from human to zebrafish; however, the amino (N) and carboxy (C) termini

are more divergent (Zhang et al., 2013). Accordingly, we hypothesize that Rpl22 and Like1 are able to bind to a largely overlapping set of RNA targets but have opposing effects on those targets, most likely because of the influence of their divergent termini. The termini might alter the way the RNA-binding domains function. Alternatively, the termini of Rpl22 and Like1 might recruit distinct *trans*-acting factors that serve as the effectors of antagonism. The preferential association of hnRNP-A1 with Rpl22 and not Like1 certainly supports this notion, but the basis for the selective interaction of hnRNP-A1 with Rpl22 remains to be determined. Nevertheless, the simplest interpretation of our data is that Like1 binding to the intronic sequences

of a pre-mRNA target interferes with the ability of Rpl22 to recruit hnRNP-A1, thereby preventing it from disrupting normal splicing. It is unclear whether Like1 alone is capable of opposing the disruptive effects of Rpl22 and hnRNP-A1 or if it requires assistance from a positive regulator of splicing. Interestingly, the recurrent G-rich element observed in the mis-spliced targets of Like1 morphants can be bound either by splicing repressors or activators (Expert-Bezançon et al., 2004). Indeed, SR proteins ASF/SF2 and SC35 can compete with hnRNP-A1 to bind to same G-rich element. hnRNP-A1 binding has been reported to antagonize the action of SR proteins and cause mis-splicing of beta-tropomyosin exon 6 (Expert-Bezançon et al., 2004). Accordingly, it is possible that Like1 may require the physical or genetic interaction with a factor(s) that facilitates normal splicing (e.g., SR proteins) in order to overcome the splice disruption induced by hnRNP-A1 and Rpl22. Our finding that some paralogs are not functionally redundant but can instead perform biologically important antagonistic functions is unusual but not without precedent. Indeed, a recent report revealed that paralogs Upf3a and Upf3b perform antagonistic functions in regulating nonsense mediated decay (Shum et al., 2016). Thus, the use of antagonistic paralogs as molecular rheostats to fine-tune biological processes appears to be an emerging theme.

Our findings reveal that Rpl22 and Like1 are not functionally redundant during early development but instead act as an antagonistic regulatory node whereby the balance of these antagonistic activities is crucial. This balance appears to be set not only by their expression level but also by control of their localization in the nucleus. Beyond their role in causing developmental abnormalities in ribosomopathies, inactivation of RP has also been associated with increased cancer risk both in inherited ribosomopathy syndromes and when RP inactivation occurs somatically. This raises the possibility that alterations in pre-mRNA splicing caused by Rpl22/Like1 imbalances might also occur in, and contribute to, transformation. We have shown that Rpl22 is a tumor suppressor as inactivation of *Rpl22* promotes lymphoma formation (Rao et al., 2012). In contrast to Rpl22, the Like1 locus (*RPL22L1*) is frequently amplified in many types of human cancers (data not shown). Moreover, mutations in RNA splicing factors (e.g., U2AF1, SF3B1, and SFPQ) have been recently implicated in the pathogenesis of human MDS and acute myelogenous leukemia (Dolnik et al., 2012; Yoshida et al., 2011). Thus, imbalances in Rpl22 and Like1 expression might disrupt splicing in these diseases when splice factors are not mutated. Understanding how Rpl22 and Like1 interact with the splicing machinery to control *smad2* splicing, may reveal insights not only into how mutations in RP can cause physical defects but also how they alter cancer risk in humans.

EXPERIMENTAL PROCEDURES

Zebrafish

Zebrafish were bred and maintained at 28.5°C under standard aquaculture conditions in the Fox Chase Association for Assessment and Accreditation of Laboratory Animal Care (AALAC)-accredited Zebrafish Facility under the auspices of an Institutional Animal Care and Use Committee (IACUC) protocol. Embryos were staged as described previously (Kimmel et al., 1995).

Morpholino Antisense Oligonucleotides

MOs (GeneTools) were designed against the ATG translational start sites of the indicated targets as previously reported (Zhang et al., 2013).

Plasmid Construction, RNA Synthesis, and Overexpression

Epitope tagged constructs were generated by standard molecular biological approaches, following which mRNA were produced by in vitro transcription and injected into one-cell embryos to achieve overexpression.

In Situ Hybridizations

Whole-mount RNA in situ hybridizations were performed as previously described (Thisse and Thisse, 2008). The stained embryos were mounted in 3% methylcellulose and photographed from the Nikon SMZ1500 stereomicroscope.

Antibodies, Western Blotting, and Immunofluorescent Staining

Immunoblotting and immunofluorescent analysis of zebrafish embryos was performed as described (Zhang et al., 2013).

Generation of *Rpl22l1*^{-/-} Mice

Rpl22l1^{-/-} mice were produced using the Targeting Vector 4595 D8 in order to insert LoxP sites in the introns between exons 1 and 2 and between exons 3 and 4. Following electroporation into R1 embryonic stem cells (ESCs), the ESCs were injected into C57BL/6 blastocysts. To ubiquitously disrupt *Rpl22l1* expression, *Rpl22l1*^{Loxp/+} mice were mated to *Mox2-cre* mice to generate *Rpl22l1*^{Loxp/-} mice, which were subsequently bred to C57BL/6 mice to create the *Rpl22l1*^{+/-} mice used in timed matings. Maintenance and analysis of these mice were performed under the auspices of an IACUC-approved protocol.

RT-PCR and Quantitative Real-Time PCR

Total RNA was isolated from embryos using Trizol (Life Technologies) and glycogen (Ambion), following which RNA was reverse transcribed and subjected to RT-PCR or quantitative real-time PCR as indicated using SYBR-Green detection.

RNA-Seq Analysis

RNA-seq analysis was performed on RNA extracted from zebrafish embryos at the end of gastrulation (10 hpf). Briefly, the mRNA-seq library was prepared from the total RNA using poly(A) selection (Truseq RNA Sample Preparation Kit V2, Illumina). RNA concentration was quantitated using a NanoDrop, and RNA integrity was measured using a BioAnalyzer chip (Agilent), followed by 50–100 bp paired end sequencing on a HiSeq2000 according to manufacturer protocols (Illumina). RNA-seq sequence-read data for control and Like1 morphants was deposited into the NCBI Short Read Archive (SRA) database (SRP093436). mRNA-seq reads for both control and Like1 morphants were mapped to the latest zebrafish genome assembly (Zv9) using the TopHat (version 2) alignment algorithm (Trapnell et al., 2009). For detection of splicing changes in Like1 morphants, the MATS algorithm was implemented using the aligned BAM files (Shen et al., 2012). Each splicing change was visualized using the IGV program (Integrative Genomics Viewer). Enrichment analysis for Gene Ontology (GO) terms was assessed using the GOstats program (Falcon and Gentleman, 2007). To evaluate the splicing strength for splice sites, we used the MaxEntScan algorithm, based on the maximum entropy model using candidate genes from Like1 morphant RNA-seq data (Yeo and Burge, 2004). The significance of differences in the strength of 5' and 3' splice sites was evaluated using the paired Wilcoxon signed rank test.

Co-immunoprecipitation of Rpl22 and Like1 with hnRNP-A1

Co-immunoprecipitation of Rpl22 and Like1 with hnRNP-A1 was performed on detergent extracts of zebrafish embryos using minor modifications to a well-established protocol (Little and Mullins, 2009).

Embryonic RNA-Protein Crosslinking and Immunoprecipitation

To develop the E-CLIP method for Rpl22/Like1 interaction with *smad2* mRNA in zebrafish embryos, we modified published protocols (Lu et al., 2013; Niranjankumari et al., 2002; Ule et al., 2003). Briefly, mRNAs encoding HA-tagged Rpl22 and Like1 were injected into one-cell embryos. Embryos were collected

at 10 hpf, crosslinked by adding 37% formaldehyde (Sigma) to 1.85% vol/vol final concentration, and lysed in hypotonic lysis buffer (10 mM Tris-HCl [pH 7.5], 10 mM NaCl, 0.5% NP40, containing protease inhibitors and SuperRNase inhibitor). Isolated nuclei were lysed by sonication (Bioruptor sonication system), following which RNA was partially trimmed by treatment with RNase T1 (Ambion) for 5 min at 37°C as described (Lu et al., 2013). The nuclear extract was clarified by centrifugation at 22,000 × g for 30 min, pre-cleared with Protein A Magnetic Beads (NEB), and immunoprecipitated with the indicated antibodies following which the immunisolated RNA was quantified as above.

Statistical Analysis

A two-tailed Student's t test was used in the experiments shown in Figures 1B, 2D, and 7B. Statistical significance was accepted at $p < 0.05$. All experiments were repeated at least three times.

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures and seven figures and can be found with this article online at <http://dx.doi.org/10.1016/j.celrep.2016.12.034>.

AUTHOR CONTRIBUTIONS

Y.Z. designed and performed all the experiments, interpreted data, and wrote the manuscript. S.P. and S.M. contributed to the RNA-seq analysis. D.J.K., M.W., J.R., and J.Z. helped with experiments. P.S.A., Q.F., and D.R.M. produced the *Rpl22l1* $-/-$ mice. M.N.O. analyzed the development of *Rpl22l1* $-/-$ mice. B.K.K. oversaw those analyses. D.L.W. oversaw the design and research, interpreted all the data, and co-wrote the manuscript with Y.Z.

ACKNOWLEDGMENTS

This work was supported by NIH grants AI110985 (D.L.W.) and P30CA006927 (D.L.W.), an appropriation from the Commonwealth of Pennsylvania (D.L.W.), and the Bishop Fund (D.L.W.). Y.Z. was supported in part by grants from the MD Anderson Cancer Center Leukemia SPORE (CA100632), When Everyone Survives Foundation, a William J. Avery postdoctoral fellowship, and an institutional postdoctoral training grant (T32 CA009035). We are grateful to the Wiest laboratory and Dr. Pengfei Xu for helpful suggestions and Drs. Siddharth Balachandran and Charles Query for critical evaluation of the manuscript. We also thank Ms. Robin G. Kunkel, MS, for illustrations. These studies were supported by the following core facilities at Fox Chase: Genomics, DNA Sequencing, Imaging, and Laboratory Animal/Zebrafish.

Received: August 6, 2016

Revised: October 15, 2016

Accepted: December 12, 2016

Published: January 3, 2017

REFERENCES

Anderson, S.J., Lauritsen, J.P., Hartman, M.G., Foushee, A.M., Lefebvre, J.M., Shinton, S.A., Gerhardt, B., Hardy, R.R., Oravec, T., and Wiest, D.L. (2007). Ablation of ribosomal protein L22 selectively impairs alphabeta T cell development by activation of a p53-dependent checkpoint. *Immunity* 26, 759–772.

Bolze, A., Mahlaoui, N., Byun, M., Turner, B., Trede, N., Ellis, S.R., Abhyankar, A., Itan, Y., Patin, E., Brebner, S., et al. (2013). Ribosomal protein SA haploinsufficiency in humans with isolated congenital asplenia. *Science* 340, 976–978.

Dick, A., Mayr, T., Bauer, H., Meier, A., and Hammerschmidt, M. (2000). Cloning and characterization of zebrafish *smad2*, *smad3* and *smad4*. *Gene* 246, 69–80.

Dolnik, A., Engemann, J.C., Scharfenberger-Schmeer, M., Mauch, J., Kelkenberg-Schade, S., Haldemann, B., Fries, T., Krönke, J., Kühn, M.W., Paschka, P., et al. (2012). Commonly altered genomic regions in acute myeloid leukemia

are enriched for somatic mutations involved in chromatin remodeling and splicing. *Blood* 120, e83–e92.

Dunn, N.R., Koonce, C.H., Anderson, D.C., Islam, A., Bikoff, E.K., and Robertson, E.J. (2005). Mice exclusively expressing the short isoform of *Smad2* develop normally and are viable and fertile. *Genes Dev.* 19, 152–163.

Expert-Bezançon, A., Sureau, A., Durosay, P., Salesse, R., Groeneveld, H., Le-caer, J.P., and Marie, J. (2004). hnRNP A1 and the SR proteins ASF/SF2 and SC35 have antagonistic functions in splicing of beta-tropomyosin exon 6B. *J. Biol. Chem.* 279, 38249–38259.

Falcon, S., and Gentleman, R. (2007). Using GOstats to test gene lists for GO term association. *Bioinformatics* 23, 257–258.

Gabut, M., Chaudhry, S., and Blencowe, B.J. (2008). SnapShot: The splicing regulatory machinery. *Cell* 133, 192.e1.

Heisenberg, C.P., and Solnica-Krezel, L. (2008). Back and forth between cell fate specification and movement during vertebrate gastrulation. *Curr. Opin. Genet. Dev.* 18, 311–316.

Heyer, J., Escalante-Alcalde, D., Lia, M., Boettinger, E., Edelmann, W., Stewart, C.L., and Kuchelapati, R. (1999). Postgastrulation *Smad2*-deficient embryos show defects in embryo turning and anterior morphogenesis. *Proc. Natl. Acad. Sci. U S A* 96, 12595–12600.

Kane, D.A., and Kimmel, C.B. (1993). The zebrafish midblastula transition. *Development* 119, 447–456.

Kearse, M.G., Ireland, J.A., Prem, S.M., Chen, A.S., and Ware, V.C. (2013). *Rpl22e*, but not *Rpl22e*-like-PA, is SUMOylated and localizes to the nucleoplasm of *Drosophila* meiotic spermatocytes. *Nucleus* 4, 241–258.

Kimelman, D., and Griffin, K.J. (2000). Vertebrate mesoderm induction and patterning. *Curr. Opin. Genet. Dev.* 10, 350–356.

Kimmel, C.B., Ballard, W.W., Kimmel, S.R., Ullmann, B., and Schilling, T.F. (1995). Stages of embryonic development of the zebrafish. *Dev. Dyn.* 203, 253–310.

Komili, S., Farny, N.G., Roth, F.P., and Silver, P.A. (2007). Functional specificity among ribosomal proteins regulates gene expression. *Cell* 131, 557–571.

Kristensen, A.R., Gsponer, J., and Foster, L.J. (2012). A high-throughput approach for measuring temporal changes in the interactome. *Nat. Methods* 9, 907–909.

Langdon, Y.G., and Mullins, M.C. (2011). Maternal and zygotic control of zebrafish dorsoventral axial patterning. *Annu. Rev. Genet.* 45, 357–377.

Little, S.C., and Mullins, M.C. (2009). Bone morphogenetic protein heterodimers assemble heteromeric type I receptor complexes to pattern the dorso-ventral axis. *Nat. Cell Biol.* 11, 637–643.

Liu, Y., Festing, M.H., Hester, M., Thompson, J.C., and Weinstein, M. (2004). Generation of novel conditional and hypomorphic alleles of the *Smad2* gene. *Genesis* 40, 118–123.

Lopez, A.J. (1998). Alternative splicing of pre-mRNA: developmental consequences and mechanisms of regulation. *Annu. Rev. Genet.* 32, 279–305.

Lu, X., Göke, J., Sachs, F., Jacques, P.E., Liang, H., Feng, B., Bourque, G., Bubulya, P.A., and Ng, H.H. (2013). SON connects the splicing-regulatory network with pluripotency in human embryonic stem cells. *Nat. Cell Biol.* 15, 1141–1152.

Matera, A.G., and Wang, Z. (2014). A day in the life of the spliceosome. *Nat. Rev. Mol. Cell Biol.* 15, 108–121.

Mukhopadhyay, R., Ray, P.S., Arif, A., Brady, A.K., Kinter, M., and Fox, P.L. (2008). DAPK-ZIPK-L13a axis constitutes a negative-feedback module regulating inflammatory gene expression. *Mol. Cell* 32, 371–382.

Narla, A., and Ebert, B.L. (2010). Ribosomopathies: human disorders of ribosome dysfunction. *Blood* 115, 3196–3205.

Niranjanakumari, S., Lasda, E., Brazas, R., and Garcia-Blanco, M.A. (2002). Reversible cross-linking combined with immunoprecipitation to study RNA-protein interactions in vivo. *Methods* 26, 182–190.

Nomura, M., and Li, E. (1998). *Smad2* role in mesoderm formation, left-right patterning and craniofacial development. *Nature* 393, 786–790.

- Rao, S., Lee, S.Y., Gutierrez, A., Perrigoue, J., Thapa, R.J., Tu, Z., Jeffers, J.R., Rhodes, M., Anderson, S., Oravec, T., et al. (2012). Inactivation of ribosomal protein L22 promotes transformation by induction of the stemness factor, Lin28B. *Blood* 120, 3764–3773.
- Schier, A.F. (2007). The maternal-zygotic transition: death and birth of RNAs. *Science* 316, 406–407.
- Schroder, P.A., and Moore, M.J. (2005). Association of ribosomal proteins with nascent transcripts in *S. cerevisiae*. *RNA* 11, 1521–1529.
- Shen, S., Park, J.W., Huang, J., Dittmar, K.A., Lu, Z.X., Zhou, Q., Carstens, R.P., and Xing, Y. (2012). MATS: a Bayesian framework for flexible detection of differential alternative splicing from RNA-seq data. *Nucleic Acids Res.* 40, e61.
- Shum, E.Y., Jones, S.H., Shao, A., Dumdie, J., Krause, M.D., Chan, W.K., Lou, C.H., Espinoza, J.L., Song, H.W., Phan, M.H., et al. (2016). The antagonistic gene paralogs Upf3a and Upf3b govern nonsense-mediated RNA decay. *Cell* 165, 382–395.
- Smith, K.A., Noël, E., Thurlings, I., Rehmann, H., Chocron, S., and Bakkers, J. (2011). Bmp and nodal independently regulate lefty1 expression to maintain unilateral nodal activity during left-right axis specification in zebrafish. *PLoS Genet.* 7, e1002289.
- Solnica-Krezel, L. (2005). Conserved patterns of cell movements during vertebrate gastrulation. *Curr. Biol.* 15, R213–R228.
- Stemple, D.L. (2000). Vertebrate development: the fast track to nodal signaling. *Curr. Biol.* 10, R843–R846.
- Takagi, M., Absalon, M.J., McLure, K.G., and Kastan, M.B. (2005). Regulation of p53 translation and induction after DNA damage by ribosomal protein L26 and nucleolin. *Cell* 123, 49–63.
- Thisse, C., and Thisse, B. (2008). High-resolution in situ hybridization to whole-mount zebrafish embryos. *Nat. Protoc.* 3, 59–69.
- Thisse, B., Wright, C.V., and Thisse, C. (2000). Activin- and Nodal-related factors control antero-posterior patterning of the zebrafish embryo. *Nature* 403, 425–428.
- Trapnell, C., Pachter, L., and Salzberg, S.L. (2009). TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 25, 1105–1111.
- Ule, J., Jensen, K.B., Ruggiu, M., Mele, A., Ule, A., and Darnell, R.B. (2003). CLIP identifies Nova-regulated RNA networks in the brain. *Science* 302, 1212–1215.
- Vautier, D., Chesné, P., Cunha, C., Calado, A., Renard, J.P., and Carmo-Fonseca, M. (2001). Transcription-dependent nucleocytoplasmic distribution of hnRNP A1 protein in early mouse embryos. *J. Cell Sci.* 114, 1521–1531.
- Waldrip, W.R., Bikoff, E.K., Hoodless, P.A., Wrana, J.L., and Robertson, E.J. (1998). Smad2 signaling in extraembryonic tissues determines anterior-posterior polarity of the early mouse embryo. *Cell* 92, 797–808.
- Wan, F., Anderson, D.E., Barnitz, R.A., Snow, A., Bidere, N., Zheng, L., Hegde, V., Lam, L.T., Staudt, L.M., Levens, D., et al. (2007). Ribosomal protein S3: a KH domain subunit in NF-kappaB complexes that mediates selective gene regulation. *Cell* 131, 927–939.
- Wang, Z., and Burge, C.B. (2008). Splicing regulation: from a parts list of regulatory elements to an integrated splicing code. *RNA* 14, 802–813.
- Wang, Z., Rolish, M.E., Yeo, G., Tung, V., Mawson, M., and Burge, C.B. (2004). Systematic identification and analysis of exonic splicing silencers. *Cell* 119, 831–845.
- Warner, J.R., and McIntosh, K.B. (2009). How common are extraribosomal functions of ribosomal proteins? *Mol. Cell* 34, 3–11.
- Xue, S., and Barna, M. (2012). Specialized ribosomes: a new frontier in gene regulation and organismal biology. *Nat. Rev. Mol. Cell Biol.* 13, 355–369.
- Yeo, G., and Burge, C.B. (2004). Maximum entropy modeling of short sequence motifs with applications to RNA splicing signals. *J. Comput. Biol.* 11, 377–394.
- Yoshida, K., Sanada, M., Shiraishi, Y., Nowak, D., Nagata, Y., Yamamoto, R., Sato, Y., Sato-Otsubo, A., Kon, A., Nagasaki, M., et al. (2011). Frequent pathway mutations of splicing machinery in myelodysplasia. *Nature* 478, 64–69.
- Zhang, Y., and Lu, H. (2009). Signaling to p53: ribosomal proteins find their way. *Cancer Cell* 16, 369–377.
- Zhang, Y., Duc, A.C., Rao, S., Sun, X.L., Bilbee, A.N., Rhodes, M., Li, Q., Kappes, D.J., Rhodes, J., and Wiest, D.L. (2013). Control of hematopoietic stem cell emergence by antagonistic functions of ribosomal protein paralogs. *Dev. Cell* 24, 411–425.
- Zhou, X., Hao, Q., Liao, J.M., Liao, P., and Lu, H. (2013). Ribosomal protein S14 negatively regulates c-Myc activity. *J. Biol. Chem.* 288, 21793–21801.
- Zhu, J., Mayeda, A., and Krainer, A.R. (2001). Exon identity established through differential antagonism between exonic splicing silencer-bound hnRNP A1 and enhancer-bound SR proteins. *Mol. Cell* 8, 1351–1361.