

Review

Cisgenesis and genome editing: combining concepts and efforts for a smarter use of genetic resources in crop breeding

TEODORO CARDI

Centro di ricerca per l'orticoltura, Consiglio per la ricerca in agricoltura e l'analisi dell'economia agraria (CREA), via Cavallegeri 25, 84098 Pontecagnano, Salerno Italy; Corresponding author, E-mail: teodoro.cardi@entecra.it

*With 1 figure and 2 tables**Received October 17, 2015 / Accepted November 27, 2015**Communicated by R. Varshney***Abstract**

Plant genetic resources (PGR) represent valuable sources of genetic variability for crop breeding. The development of novel biotechnologies is necessary for increasing the efficiency of their use in pre-breeding and breeding work. The genome sequencing of hundreds of genotypes and the mining of allele diversity in major crops and populations of landraces and wild relatives allow the isolation of genes underlying characters of interest and their precise modification or transfer into targeted varieties. The technological developments and applications of new plant breeding techniques (NPBT) that maximize the similarity with gene transfer by crossing (cisgenesis/intragenesis) or the accuracy of biotechnological approaches (genome editing) are reviewed. Their potentialities and current limitations as well as the possible advantages of using them separately or combined for the exploitation of PGR in crop breeding are also discussed. Above-mentioned NPBT tackle some objections to the application of biotechnologies in agriculture and are under review worldwide to assess the possible exclusion from the current regulation systems for genetically modified plants.

Key words: cisgenesis — intragenesis — genome editing — plant genetic resources — crop breeding — nucleases

Plant genetic resources (PGR), including landraces (LR), heirloom varieties (HV) and crop wild relatives (CWR), are paramount important sources of genetic variability at the intraspecific and interspecific level for breeding new crop varieties. Due to their history, based on natural and artificial (farmer) selection in particular environments, LRs are adapted to local growing conditions and show interesting features in terms of resilience and organoleptic and nutritional value (Villa et al. 2005, Zeven 1998). Besides their indirect use as a source of genes for the development of new varieties, they can be also used directly by farmers. Nevertheless, they often show some specific defects, for example susceptibility to one or more pathogen strains, that limit their use. Although LRs and CWRs are being used in intraspecific/interspecific hybridizations and introgression breeding for long time (Andersen et al. 2015, Jacobsen and Schouten 2007, Palmgren et al. 2015), the development of novel technologies is necessary for increasing the efficiency of their use in pre-breeding and breeding work, in order to promptly respond to present and future agricultural challenges.

At the end of the last century, genetic engineering techniques enabled the production of the first generation of genetically modified plants (GMP or transgenic plants), based on the transfer and random insertion into the host plant genomes of genes isolated from other plant species or from other organisms. In this century, the improvement of sequencing technologies has allowed the complete deciphering of genomes in many species

of agricultural interest, such as rice, maize, tomato, potato, pepper, eggplant, cucumber, melon and others (Bolger et al. 2014, Hamilton and Buell 2012, Michael and VanBuren 2015). Next-generation sequencing (NGS) technologies consent the whole- or targeted genome resequencing of hundreds of genotypes and the mining of allele diversity in the populations of crop landraces and wild relatives (Causse et al. 2013, Lin et al. 2014, Mascher et al. 2013, Neves et al. 2013, Sainetnac et al. 2011). Knowledge of the structure and function of plant genomes in major agricultural and related PGR prompts the development of so-called second-generation biotechnologies and their application in breeding (new plant breeding techniques or NPBT), aiming to the isolation of genes underlying the characters of interest and their precise modification or transfer into targeted varieties (Gruskin 2012, He et al. 2014, Lusser et al. 2012, Varshney et al. 2014).

In relation to the use of transgenic plants in agriculture, the transfer in cultivated species of genes isolated from other species and/or the random insertion of the transgene into the genome of crop plants are sometimes perceived as potential risks to the environment and health. NPBT that maximize the similarity with gene transfer by crossing (cisgenesis/intragenesis) or the accuracy of biotechnological approaches (genome editing) tackle some objections to the application of biotechnologies in agriculture. They are under review worldwide to assess the possible exclusion from the current GMP regulation systems (Gruskin 2012, Hunter 2014, Kuzma and Kokotovich 2011, Podevin et al. 2012). Hereinafter, the recent developments of cisgenic/intragenic and genome editing approaches based on the advancement of sequencing technologies and the availability of genome information are reviewed, evaluating their potentialities and current limitations as well as the possible advantages of using them separately or combined for the exploitation of PGR in crop breeding.

Cisgenesis/Intragenesis

Although several variations have been proposed (Holme et al. 2013), both cisgenic and intragenic approaches are based on the transfer of only genes and regulatory sequences derived from other genotypes of the same or sexually compatible species. In case of cisgenesis, the entire gene with its own regulatory sequences and in sense orientation is transferred. In the second case (intragenesis), different coding and regulatory sequences are assembled either in sense or in antisense orientation, the latter if the aim is to reduce gene expression by activating the RNA

interference (RNAi) pathway. Further, according to some authors, P-DNA borders and vector backbone sequences derived from the sexually compatible gene pool should be used in intragenic vectors for *Agrobacterium*-mediated transformation (Conner et al. 2007). While, at least in principle, cisgenic products can be achieved by conventional breeding, the same is not true for intragenic ones. Anyway, in both approaches, only the genes that code for the characters to be modified should be eventually present in regenerated plants and a number of technologies can be adopted to avoid the presence of selectable marker genes generally used for *in vitro* selection of transformed cells (Holme et al. 2013 for review). Conventional gene delivery methods, such as either *Agrobacterium* spp. or biolistics-based transformation, are commonly used for transformation.

As recently reviewed (Espinoza et al. 2013, Holme et al. 2013, Lamalakshmi Devi et al. 2013), the cisgenic approach has been used to improve pathogen resistance and quality traits in several crops, namely durum wheat, poplar, grape, apple, potato. Approaches aimed at increasing intragenic gene expression have been published in apple, strawberry and potato to increase the resistance to pathogens and in perennial ryegrass tolerance to drought, whereas the targeted reduction in gene expression was obtained in potato and alfalfa with the aim to improve various aspects of quality (Table 1). Not all products so far classified as intragenic or cisgenic, however, fully comply with requirements set for such kinds of plants, because some still contain microbial regulatory sequences and/or selectable marker genes (Holme et al. 2013). More recently, molecular characterization of true cisgenic apple plants previously produced and expressing the *Rvi6* scab resistance genes has been reported in detail (Vanblaere et al. 2014), while new cisgenic apples with the same trait have been developed using an alternative recombinase system (Würdig et al. 2015). Furthermore, marker-free cisgenic potato (*Solanum tuberosum*) plants expressing late blight resistance genes from *S. stoloniferum* (*Rpi-sto1*) and *S. venturii* (*Rpi-vnt1.1*) have been produced by *Agrobacterium*-mediated transformation without using any marker gene but PCR for the selection of transformed plants. Due to the activity of both introduced *R* genes, cisgenic plants showing broad-spectrum late blight resistance could be selected (Jo et al. 2014). An intragenic vector for future applications has been developed in *Citrus* spp. (An et al. 2013).

Because the sequences transferred by either cisgenesis or intragenesis are derived from the same or related species, the knowledge of their sequence, position and function in the genomes of origin is essential. It is expected that such knowledge will

prompt a wider use of both technologies in place of common transgenesis. Although gene transfer within the same or evolutionarily close species can also be achieved by conventional breeding, cisgenic/intragenic approaches reduce considerably both the duration of selection steps and linkage drag. In addition, the genotype and phenotype of varieties to be modified remain largely unchanged, an issue particularly important with vegetatively propagated heterozygous and/or long-cycle species, such as potato or, in general, fruit trees. Another potential advantage of cisgenesis/intragenesis, compared to conventional breeding, is the higher knowledge of transferred sequences (Conner et al. 2007, Holme et al. 2013).

The EFSA (European Food Safety Authority) concluded that cisgenic plants pose risks similar to those obtained with conventional breeding, while the risks of plants produced by intragenesis are similar to those of transgenic plants (EFSA Panel on Genetically Modified Organisms (GMO), 2012b). Furthermore, several recent reports confirmed a greater acceptance by consumers of cisgenic products compared to the corresponding transgenics (Delwaide et al. 2015, Holme et al. 2013). A number of field trials with either cisgenic or intragenic plants are being carried out worldwide and the applications for deregulation are pending (Holme et al. 2013). As far as the regulation of these products is concerned, generally speaking, United States, Canada and Australia are showing a more open orientation and are aiming to distinguish them from conventional GMPs (Hunter 2014, Lusser and Davies 2013), while in Europe and other countries, the attitude is more cautious.

Notwithstanding the aforementioned considerations, cisgenic/intragenic approaches still show some drawbacks that limit a wider application. A careful analysis of regulatory sequence type and size used in cisgenic/intragenic constructs is necessary. In apple, although an acceptable level of scab resistance was achieved, differences in expression level of the *Rvi6* gene depended on using the *rbcS* promoter from the same species or the native promoter with different lengths (Joshi et al. 2011, Vanblaere et al. 2014, Würdig et al. 2015). Additional effects of the terminator type and length were discussed by the same authors. Variability in gene expression could be also due to random insertion of the cisgene in the host genome, determining negative position-dependent epigenetic regulation (Vanblaere et al. 2014). Further, random integration of cisgenes/intragenes can potentially determine, similar to regular transgenes, the interruption and silencing of resident genes or other relevant sequences and so a thorough screening of regenerated plants is

Table 1: Examples of agronomic traits modified through the application of either *cisgenesis* or *intragenesis* in various crops

Species	Trait	Gene	Technology	References
Alfalfa	Lignin content	<i>Comt</i>	Intragenesis	Weeks et al. (2008)
Apple	Scab resistance	<i>Rvi6</i> (<i>HcrVf2</i>)	Cisgenesis, intragenesis	Joshi et al. (2011); Vanblaere et al. (2014); Würdig et al. (2015)
Barley	Grain phytase activity	<i>HvPAPhy_a</i>	Cisgenesis	Holme et al. (2012)
Durum wheat	Baking quality	<i>1Dy10</i>	Cisgenesis	Gadaleta et al. (2008)
Perennial ryegrass	Drought tolerance	<i>Lvpv1</i>	Intragenesis	Bajaj et al. (2008)
Poplar	Plant growth and stature, wood properties	<i>PtGA20ox7</i> , <i>PtGA2ox2</i> , <i>PtRGL1_1</i> , <i>PtRGL1_2</i> , <i>PtGAIL</i>	Cisgenesis	Han et al. (2011)
Potato	Late blight resistance	<i>Rpi-sto1</i> , <i>Rpi-vnt1.1</i>	Cisgenesis	Jo et al. (2014)
	High amylopectin	<i>GBSS</i>	Intragenesis	de Vetten et al. (2003)
	Prevention of black spot bruise	<i>Ppo</i>	Intragenesis	Rommens et al. (2004)
	Accumulation of reducing sugars after cold storage and acrylamide after high-temperature processing	<i>R1</i> , <i>PhL</i> , <i>StAs1</i> , <i>StAS2</i>	Intragenesis	Chawla et al. (2012); Rommens et al. (2006, 2008)
Strawberry	Grey mould resistance	<i>PGIP</i>	Intragenesis	Schaart (2004)

necessary. Finally, the number of gene copies and the amount of vector backbone sequences transferred is an additional issue, because up to 80% of original plants regenerated from cisgenic transformation experiments showed the integration of vector backbone sequences (Jo et al. 2014, Vanblaere et al. 2014, 2011). New vectors containing buffer plant sequences or genes for counterselection in the backbone can help limit the negative consequences of unprecise integration of cisgenes and intragenes in the host genome. Furthermore, vectors composed of only plant-derived sequences have been proposed. The use of minimal linear cassettes and biolistic gene delivery is also a valid method to have 'cleaner' transformations (An et al. 2013, Conner et al. 2007, Gadaleta et al. 2008, Holme et al. 2013).

Genome Editing

'Genome editing' includes a set of techniques that allow to edit, delete, replace or insert, in a targeted site, specific genomic sequences of interest. They are based on the induction, in different organisms, including plants, of cuts in double-strand DNA (DSB, double-strand breaks), which are then 'repaired' with two different processes: the non-homologous end-joining (NHEJ) or homology-directed repair (HDR) (Gaj et al. 2013, Puchta and Fauser 2014, Rinaldo and Ayliffe 2015, Voytas 2013).

The breaks in the double-strand DNA can be induced by four systems based on specific enzymes: (i) meganucleases, (ii) zinc finger nucleases (ZFN), (iii) transcription activator-like effector nucleases (TALEN) and (iv) clustered regular interspaced short palindromic repeats/CRISPR-associated nucleases (CRISPR/Cas). In comparison with the meganucleases, the other systems allow to edit more efficiently 'target' sequences and hence have a larger use than the former for different purposes. ZFN and TALEN are based on the association between some proteins, which have the property to recognize and bind in a selective manner to pre-selected nucleotide sequences, and two units of a nuclease (*FokI*), which, when dimerized, cut double-strand DNA. In order to obtain dimerization of *FokI*, a couple of ZF or TALE proteins are designed and synthesized according to the nucleotide sequence to be modified. Each ZF protein recognizes three nucleotides on each DNA strand, and generally, each ZFN uses 3–4 proteins per filament: in this way, the specificity is determined by 18–24 nucleotides (9–12 per filament). In the TALEN system, the specific recognition with the DNA sequence is determined by the TALE proteins (discovered in phytopathogenic *Xanthomonas* spp. bacteria), each of which recognizes a single nucleotide. Specific recognition of the region to be modified is determined by thirty–forty repeats (15–20 per filament), each containing 33–35 amino acids. The sequence of each repeat is highly conserved, except at positions 12 and 13 (RVDs, repeat variable diresidues). Each repeat recognizes a single base. The dimeric *FokI* cuts the DNA between the 'batteries' of ZF or TALE proteins. Unlike ZFN and TALEN, in CRISPR/Cas9 technology, recognition of the DNA sequence to be modified is determined not by proteins, but by a chimeric sequence of RNA (single guide RNA – sgRNA), which results from the fusion of the two sequences (crRNA and tracrRNA) present in the natural system (the CRISPR/Cas system is used by bacteria to defend themselves against phages), while the cutting of DNA sequence is performed by the associated monomeric enzyme Cas9 (Boch et al. 2009, Christian et al. 2010, Jinek et al. 2012, Kim and Chandrasegaran 1994, Kim et al. 1996).

After the two strands of the double helix of DNA are cut, in the absence of foreign donor sequences, the filaments are

predominantly rejoined in plants by NHEJ, but, because this process is susceptible to errors, small changes (mostly frame-shift mutations due to insertions and deletions) in the original sequence are induced, which generally result in the loss of function of the target gene and a mutated phenotype. This result is similar to that obtainable with other technologies in use since long time, such as mutagenesis with chemical or physical mutagens, although the induction of mutations is not random in the genome, as in classical mutagenesis, but limited to genes of interest. As a consequence of conventional random mutagenesis, unwanted mutations can be induced throughout the genome and large-scale screens of mutagenized populations are needed to identify those plants with mutations of interest. Antisense and RNAi-based technologies can be also used to selectively knock-down gene expression. Their effects, however, are often incomplete, not stable across generations and not limited to the gene of interest. In addition, they target transcripts, but no other genetic elements, such as promoters, enhancers, introns and intergenic regions, which are instead accessible by genome editing tools. If, together with the nucleases described above, appropriate DNA fragments homologous to the target sequence are also inserted into the cell, they can, using the mechanism of homologous recombination (HDR), which unlike NHEJ does not induce errors, replace (correct) some nucleotide sequences of the gene to be modified or add new genes or regulatory elements in a predetermined position of the genome (Chen and Gao 2014). The induction of site-specific random mutations (1), the induction of mutations in a predefined sequence of a particular gene (2) and the replacement or the insertion of an entire gene (3) are collectively dubbed as SDN (site-directed nuclease)-1, SDN-2 and SDN-3, respectively (EFSA Panel on Genetically Modified Organisms (GMO), 2012a).

Delivery into plant cells of various components necessary for genome editing has been accomplished by either transient or stable transformation methods, including protoplast systems, *Agrobacterium*-based procedures, biolistics and virus vectors. Transient methods are preferable both for the cytotoxicity sometimes shown when nucleases are expressed stably and for regulatory reasons, in order to avoid the presence of stably integrated exogenous sequences. The choice of one method or another, however, depends on several factors, such as the length of life cycle, the feasibility of protoplast isolation and plant regeneration, the type of molecule to be expressed, the repairing mechanism and the editing objective pursued (Baltes and Voytas 2015, Baltes et al. 2014, Belhaj et al. 2013, 2015, Bortesi and Fischer 2015, Fichtner et al. 2014, Kumar and Jain 2015, Schaeffer and Nakata 2015, Voytas 2013).

The three main methods presently available for genome editing in plants have been recently compared for several aspects in various reviews (Belhaj et al. 2015, Bortesi and Fischer 2015, Chen and Gao 2014, Fichtner et al. 2014, Kumar and Jain 2015, Liu and Fan 2014, Mahfouz et al. 2014). In comparison with ZFNs and TALENs, the CRISPR/Cas approach looks more attractive based on higher simplicity, accessibility, cost, versatility, possibility of multiplexing and other aspects, including an easier open access to resources. Although contrasting results have been reported, the off-target effects remain the main concern for a wider application of genome editing to functional genomics and plant breeding. Bioinformatic analysis of target genomes is necessary to avoid targeting sequences repeated throughout the genome, and to this end, several tools available online are being developed. Furthermore, the double nickase and Cas9-nuclease fusion systems were developed to reduce

off-target editing of CRISPR/Cas9 (Belhaj et al. 2015, Schaeffer and Nakata 2015). The relatively large size of Cas9 enzyme can be a limiting factor in some cases, for example when it has to be delivered through viral vectors, whose cell-to-cell movement necessary to *in vivo* edit meristematic or gametic cells is prevented by their genome size (Schaeffer and Nakata 2015). Anyway, a novel smaller version of Cas9 isolated from *Staphylococcus aureus* edited the mouse genome with efficiencies similar to those of the widely used form from *Streptococcus pyogenes* (Ran et al. 2015). Additional future technological improvements will derive by the recent discovery in diverse bacteria of the single RNA-guided endonuclease Cpf1. In comparison with Cas9, it shows distinct advantages, including the production of sticky ends after cutting DNA instead of blunt ends, a feature that should make editing by sequence insertion/replacement easier and more controllable (Zetsche et al. 2015).

Due to difficulties in transferring in plant cells a DNA repair template together with nucleases, and to the intrinsic low efficiency of homologous recombination in higher plant nuclear genomes, most reports deal with targeted mutagenesis (gene-knockout or SDN-1 approaches). As far as crop plants are concerned, mutations in targeted genes were obtained for quality-related and pathogen or herbicide resistance traits in cereals (barley, maize, rice, sorghum and wheat) (Char et al. 2015, Gurushidze et al. 2014, Jiang et al. 2013, Li et al. 2012, Liang et al. 2014, Shan et al. 2015, Svitashv et al. 2015, Upadhyay et al. 2013, Wang et al. 2014, Wendt et al. 2013, Zhang et al. 2014), soybean (Curtin et al. 2011, Haun et al. 2014, Jacobs et al. 2015, Li et al. 2015, Sun et al. 2015), potato (Clasen et al. 2015, Nicolia et al. 2015, Wang et al. 2015b), tomato (Brooks et al. 2014, Lor et al. 2014), tobacco (Gao et al. 2015) and perennial fruit trees (poplar, sweet orange, apple and fig) (Fan et al. 2015, Jia and Wang 2014, Peer et al. 2015, Zhou et al. 2015b), although in some cases mutations were only induced in reporter genes and/or studied *in vitro* at the molecular level.

Some examples in which traits of agronomic interest were investigated in full plants are summarized in Table 2. Interestingly, they include worldwide important crops, such as potato, soybean, rice and wheat. In the latter, by TALEN, it was possible to induce mutations in all three *TaMLO* (mildew-resistance locus) homoeologs, thus conferring resistance to *Blumeria graminis* f. sp. *tritici*, a trait not found in natural populations. HDR-mediated approaches including gene editing (SDN-2) or gene replacement/stacking (SDN-3) were achieved in tobacco, soybean, barley, rice and maize crops, besides model *A. thaliana* and *Nicotiana benthamiana*. In tobacco, mutations known to confer herbicide resistance were introduced into *SuR* genes by ZFN, confirming the acquisition of resistance in calli (Townsend et al. 2009), and by TALEN (Zhang et al. 2013). *ALS* genes, conferring herbicide resistance, were successfully edited also in maize and soybean by CRISPR/Cas9 (Li et al. 2015, Svitashv et al. 2015). In barley, a single amino acid exchange inducing GFP to YFP conversion was achieved by TALEN-induced modification of a *gfp* transgene (Budhagatapalli et al. 2015), while preselected sequence changes were induced in rice *PDS*- and *EPSPS*-coding genes by CRISPR/Cas and TALEN systems, respectively (Shan et al. 2013, Wang et al. 2015a). In maize, using the ZFN technology, the endogenous *IPK1* gene, involved in phytate accumulation, was disrupted by the insertion of the *PAT* gene, encoding a phosphinothricin acetyltransferase, achieving in modified plants, at the same time, herbicide tolerance and low accumulation of phytate (Shukla et al. 2009), while, more recently, the *PAT* gene was precisely inserted at a preselected locus by HDR mediated by CRISPR/Cas9 (Svitashv et al. 2015). These studies showed that with the use of TALEN and CRISPR, the frequency of gene targeting by HDR is higher and now practicable for crop breeding.

Provided that sequences encoding nucleases and other components are not present in the final edited products, plants modified by either SDN-1 or SDN-2 approaches are indistinguishable

Table 2: Examples of agronomic traits modified through the application of *genome editing* approaches in various crops

Crop	Trait	Gene	Technology	References
Gene knockout (SDN-1)¹				
Rice	Resistance to bacterial blight	<i>OsSWEET14; OsSWEET13</i>	TALEN; CRISPR/Cas9	Li et al. (2012); Zhou et al. (2015a)
Bread wheat	Fragrance	<i>OsBADH2</i>	TALEN	Shan et al. (2015)
Maize	Resistance to powdery mildew	<i>TaMLO-A1, TaMLO-B1, TaMLO-D1</i>	TALEN	Wang et al. (2014)
	Phytate biosynthesis	<i>IPK1</i>	ZFN	Shukla et al. (2009)
	Leaf epicuticular wax composition	<i>glossy2 (gl2)</i>	TALEN	Char et al. (2015)
	Leaf development; Male fertility; Herbicide resistance	<i>LIG1; Ms26, Ms45; ALS1, ALS2</i>	CRISPR/Cas9	Svitashv et al. (2015)
Soybean	Profile and unsaturation level of seed fatty acids	<i>FAD2-1A and FAD2-1B</i>	TALEN	Haun et al. (2014)
Poplar	Lignin content; Condensed tannin content	<i>4CL1; 4CL2</i>	CRISPR/Cas9	Zhou et al. (2015b)
Potato	Accumulation of reducing sugars after cold storage and acrylamide after high-temperature processing	<i>VInv</i>	TALEN	Clasen et al. (2015)
Tomato	Accumulation of steroidal glycoalkaloids	<i>SrSSR2</i>	TALEN	Sawai et al. (2014)
	Plant development	<i>PROCERA (PRO)</i>	TALEN	Lor et al. (2014)
	Leaf development	<i>ARGONAUTE7 (SIAGO7)</i>	CRISPR/Cas9	Brooks et al. (2014)
Gene editing (SDN-2)				
Maize	Herbicide resistance	<i>ALS2</i>	CRISPR/Cas9	Svitashv et al. (2015)
Soybean	Herbicide resistance	<i>ALS1</i>	CRISPR/Cas9	Li et al. (2015)
Tobacco	Herbicide resistance ²	<i>ALS SuRA and SuRB</i>	ZFN	Townsend et al. (2009)
Gene replacement/stacking (SDN-3)				
Maize	Phytate production/herbicide resistance	<i>IPK1/PAT</i>	ZFN	Shukla et al. (2009)
	Herbicide resistance ²	<i>PAT</i>	CRISPR/Cas9	Svitashv et al. (2015)

¹SDN, site-directed nuclease.

²Tested *in vitro*.

from similar plants obtained by conventional mutagenesis (although background mutations are by far less) or natural variation. In addition, it is difficult to evidence the modification occurred in a few nucleotides and thus distinguish them from unmodified original genotypes. Hence, many argue that such plants should not be considered transgenic and thus they should fall outside the boundaries of GMP regulation (Belhaj et al. 2013, Bortesi and Fischer 2015, Chen and Gao 2014, Fichtner et al. 2014, Rinaldo and Ayliffe 2015, Sprink et al. 2015). Recent results about the possibility to carry out DNA-free editing of plant genomes with preassembled CRISPR-Cas9 ribonucleoproteins could contribute to alleviate regulatory concerns towards NPBT (Woo et al. 2015). The US Ministry of Agriculture (USDA) recently declared that the products derived from genome editing approaches will be evaluated case by case, deregulating then corn plants with low phytate obtained by the use of ZFN-3 (Jones 2015, Lusser and Davies 2013, Voytas and Gao 2014). The European Commission has not yet delivered its opinion. Only about the SDN-3 approaches, EFSA concluded that, similar to conventional transgenic approaches, risks are related to the sequence of transferred genes, although there are less risks associated with the potential disruption of either coding or regulatory sequences following the transgene insertion (EFSA Panel on Genetically Modified Organisms (GMO), 2012a). According to the same source, a reduced amount of specific data should be asked in some cases, for example, after site-specific transfer of a cisgene. Novel regulatory schemes for products obtained by different genome editing approaches have been recently proposed (Araki and Ishii 2015, Wolt et al. 2015).

Discussion and Perspectives

It is estimated that 7–15 million euro and 4–6 years are needed to obtain the authorization for the cultivation of GMP in the environment, an investment that only large multinational companies can bear (Hartung and Schiemann 2014, Jones 2015, Voytas and Gao 2014). Although in 2014 the use of GMP in agriculture reached 181.5 million hectares in the world (James 2014) and quite a number of edible crops showing biotic or abiotic stress tolerance and improved nutritional value are in the pipeline (Ricroch and Henard-Damave 2015), the scale of investment for their approval has limited the number to a few crops (soybean,

maize, cotton, canola) and a few characters (almost exclusively resistance to insects and herbicides).

A more widespread implementation of novel biotechnologies in the characterization and exploitation of plant genetic resources for crop breeding is desirable. The implementation of NPBT in breeding and organic farming has been recently advocated in order to reintroduce in crop varieties the properties present in wild relatives ('rewilding') and close the productivity gap between organic and conventional agriculture (Andersen et al. 2015, Palmgren et al. 2015). NPBT can be used in functional and complementation studies with the aim to identify major genes and superior mutations/alleles present in PGR (Fig. 1). Besides agronomic aspects, the direct use of landraces and traditional varieties is appreciated for their 'cultural' value and their link to local 'traditional' agriculture. For a number of reasons, there is a general willingness to not modify their original genotype and phenotype. Hence, SDN-1 and SDN-2 approaches, based on the use of various nucleases, are the most convenient methods to precisely correct their defects, either inhibiting the expression of specific genes or enabling novel limiting features, leaving the genetic background and other traits largely untouched. For indirect uses of landraces, traditional varieties and crop wild relatives in plant breeding, structural and functional information derived from genomic studies in collections of such PGR can be adopted to edit gene sequences in elite crop varieties, targeting specific sites and inducing relevant mutations. Finally, PGR belonging to the 'breeders' gene pool can be used also as source of cisgenes to be transferred in crop varieties, replacing alleles with superior counterparts or introducing new genes in the cultivated gene pool. Although no scientific evidence has been related so far to a safer use of cisgenes instead of transgenes, a positive consumers' perception linked to the use of genes from compatible gene pools has been reported (DeFrancesco 2013, Delwaide et al. 2015, Holme et al. 2013). Nevertheless, the unpredictability of cisgene/intragenic integration and expression is not only a limitation from the technical point of view, but also for public acceptance and regulatory issues (Schubert and Williams 2006). Hence, the combination of cisgene/intragenic concepts with those of new genome editing techniques can help reduce the concerns associated with the use of biotechnologies in agriculture, opening new perspectives for a more efficient use of genetic resources in crop breeding, valorizing the




Fig. 1: Implementation of genome sequence information and novel biotechnological approaches for the exploitation of genetic resources in crop breeding. LR: landraces; HV: heirloom varieties; CWR: crop wild relatives; SDN-1, 2, 3: site-directed nuclease usage for gene knockout, gene editing or gene replacement/stacking, respectively

genomic and phenomic data that national and international initiatives are producing (<http://www.divseek.org/>). Further, editing of genes coding for proteins involved in chromosome segregation (e.g. kinetochore protein CENH3) could facilitate the production of haploids and the fixation of useful traits derived by introgression breeding in crops where efficient methods for haploidization are not available (Chan 2010).

In order to stimulate the adoption of biotechnological innovations also by small- and medium-sized companies and public research institutes, and their use in crops of minor economic importance worldwide, many researchers believe that it would be more appropriate to adopt a procedure assessment based on the product and not on the process, with evaluations conducted case by case, excluding the products of the new technologies described in this article from the regulation of transgenic plants so far produced (Conner et al. 2007, Hartung and Schiemann 2014, Jacobsen and Schouten 2009, Jones 2015, Schouten 2014, Schouten et al. 2006, Voytas and Gao 2014). That discussion is having conflicting outcomes in different countries, reflecting the general approach adopted in the regulatory procedures: based on the assessment of the technology, as is the case in Europe with the Directive 2001/18/EC, or the product, as it happens in North America (Lusser and Davies 2013). Anyway, the potential benefits and disadvantages or risks of a larger use of biotechnologies for agriculture, health and the environment can be nowadays assessed more efficiently than in the past, sequencing the genome of modified plants and thoroughly profiling their metabolome (Ladics et al. 2015). In addition, the potential effects of the genetic modifications induced in crops can be analysed with predictive models based on ‘(Crop) Systems Biology’ approaches (Caramante et al. 2014, Keurentjes et al. 2013, 2011).

Acknowledgements

The research was conducted with the support of the ‘GenHort’ project (‘Adding value to elite Campania horticultural crops by advanced genomic technologies’, MIUR PON02_00395_3215002). The article reflects the personal views of the author and not necessarily those of the Council for Agricultural Research and Economy (CREA).

References

An, C., V. Orbović, and Z. Mou, 2013: An efficient intragenic vector for generating intragenic and cisgenic plants in citrus. *Am. J. Plant Sci.* **04**, 2131–2137.

Andersen, M. M., X. Landes, W. Xiang, A. Anyshchenko, J. Falhof, J. T. Osterberg, L. I. Olsen, A. K. Edenbrandt, S. E. Vedel, B. J. Thorsen, P. Sandoe, C. Gamborg, K. Kappel, and M. G. Palmgren, 2015: Feasibility of new breeding techniques for organic farming. *Trends Plant Sci.* **20**, 426–434.

Araki, M., and T. Ishii, 2015: Towards social acceptance of plant breeding by genome editing. *Trends Plant Sci.* **20**, 145–149.

Bajaj, S., S. Puthigae, K. Templeton, C. Bryant, G. Gill, P. Lomba, H. Zhang, F. Altpeter, and Z. Hanley, 2008: Towards engineering drought tolerance in perennial ryegrass using its own genome 6th Canadian plant genomics workshop, Abstracts, 62.

Baltes, N. J., and D. F. Voytas, 2015: Enabling plant synthetic biology through genome engineering. *Trends Biotechnol.* **33**, 120–131.

Baltes, N. J., J. Gil-Humanes, T. Cermak, P. A. Atkins, and D. F. Voytas, 2014: DNA replicons for plant genome engineering. *Plant Cell* **26**, 151–163.

Belhaj, K., A. Chaparro-Garcia, S. Kamoun, and V. Nekrasov, 2013: Plant genome editing made easy: targeted mutagenesis in model and crop plants using the CRISPR/Cas system. *Plant Methods* **9**, 39.

Belhaj, K., A. Chaparro-Garcia, S. Kamoun, N. J. Patron, and V. Nekrasov, 2015: Editing plant genomes with CRISPR/Cas9. *Curr. Opin. Biotechnol.* **32**, 76–84.

Boch, J., H. Scholze, S. Schornack, A. Landgraf, S. Hahn, S. Kay, T. Lahaye, A. Nickstadt, and U. Bonas, 2009: Breaking the code of DNA binding specificity of TAL-type III effectors. *Science* **326**, 1509–1512.

Bolger, M. E., B. Weisshaar, U. Scholz, N. Stein, B. Usadel, and K. F. Mayer, 2014: Plant genome sequencing – applications for crop improvement. *Curr. Opin. Biotechnol.* **26**, 31–37.

Bortesi, L., and R. Fischer, 2015: The CRISPR/Cas9 system for plant genome editing and beyond. *Biotechnol. Adv.* **33**, 41–52.

Brooks, C., V. Nekrasov, Z. B. Lippman, and J. Van Eck, 2014: Efficient gene editing in tomato in the first generation using the clustered regularly interspaced short palindromic repeats/CRISPR-associated9 system. *Plant Physiol.* **166**, 1292–1297.

Budhagatapalli, N., T. Rutten, M. Gurushidze, J. Kumlehn, and G. Hensel, 2015: Targeted modification of gene function exploiting homology-directed repair of TALEN-mediated double-strand breaks in barley. *G3: Genes - Genomes - Genetics* **5**, 1857–1863.

Caramante, M., N. D’Agostino, A. Venezia, and T. Cardi, 2014: Dai geni alle colture: analisi multi-livello con strategie di ‘(Crop) Systems Biology’ per un approccio interdisciplinare alla ricerca in orticoltura. *Italus Hortus* **21**, 43–56.

Causse, M., N. Desplat, L. Pascual, M. C. Le Paslier, C. Sauvage, G. Bauchet, A. Berard, R. Bounon, M. Tchoumakov, D. Brunel, and J. P. Bouchet, 2013: Whole genome resequencing in tomato reveals variation associated with introgression and breeding events. *BMC Genom.* **14**, 791.

Chan, S. W., 2010: Chromosome engineering: power tools for plant genetics. *Trends Biotechnol.* **28**, 605–610.

Char, S. N., E. Unger-Wallace, B. Frame, S. A. Briggs, M. Main, M. H. Spalding, E. Vollbrecht, K. Wang, and B. Yang, 2015: Heritable site-specific mutagenesis using TALENs in maize. *Plant Biotechnol. J.* **13**, 1002–1010.

Chawla, R., R. Shakya, and C. M. Rommens, 2012: Tuber-specific silencing of asparagine synthetase-1 reduces the acrylamide-forming potential of potatoes grown in the field without affecting tuber shape and yield. *Plant Biotechnol. J.* **10**, 913–924.

Chen, K., and C. Gao, 2014: Targeted genome modification technologies and their applications in crop improvements. *Plant Cell Rep.* **33**, 575–583.

Christian, M., T. Cermak, E. L. Doyle, C. Schmidt, F. Zhang, A. Hummel, A. J. Bogdanove, and D. F. Voytas, 2010: Targeting DNA double-strand breaks with TAL effector nucleases. *Genetics* **186**, 757–761.

Clasen, B.M., T.J. Stoddard, S. Luo, Z.L. Demorest, J. Li, F. Cedrone, R. Tibebu, S. Davison, E.E. Ray, A. Daulhac, A. Coffman, A. Yabandith, A. Retterath, W. Haun, N.J. Baltes, L. Mathis, D.F. Voytas, and F. Zhang, 2015: Improving cold storage and processing traits in potato through targeted gene knockout. *Plant Biotechnol. J.* doi: 10.1111/pbi.12370.

Conner, A., P. Barrell, S. Baldwin, A. Lokerse, P. Cooper, A. Erasmussen, J.-P. Nap, and J. E. Jacobs, 2007: Intragenic vectors for gene transfer without foreign DNA. *Euphytica* **154**, 341–353.

Curtin, S. J., F. Zhang, J. D. Sander, W. J. Haun, C. Starker, N. J. Baltes, D. Reyon, E. J. Dahlborg, M. J. Goodwin, A. P. Coffman, D. Dobbs, J. K. Joung, D. F. Voytas, and R. M. Stupar, 2011: Targeted mutagenesis of duplicated genes in soybean with zinc-finger nucleases. *Plant Physiol.* **156**, 466–473.

DeFrancesco, L., 2013: How safe does transgenic food need to be? *Nat. Biotechnol.* **31**, 794–802.

Delwaide, A. C., L. L. Nalley, B. L. Dixon, D. M. Danforth, R. M. Nayga Jr, E. J. Van Loo, and W. Verbeke, 2015: Revisiting GMOs: are there differences in European consumers’ acceptance and valuation for cisgenically vs transgenically bred rice? *PLoS ONE* **10**, e0126060.

EFSA Panel on Genetically Modified Organisms (GMO), 2012a: Scientific opinion addressing the safety assessment of plants developed

- using Zinc Finger Nuclease 3 and other Site-Directed Nucleases with similar function. *EFSA J.* **10**, 2943.
- EFSA Panel on Genetically Modified Organisms (GMO), 2012b: Scientific opinion addressing the safety assessment of plants developed through cisgenesis and intragenesis. *EFSA J.* **10**, 2561.
- Espinoza, C., R. Schlechter, D. Herrera, E. Torres, A. Serrano, C. Medina, and P. Arce-Johnson, 2013: Cisgenesis and intragenesis: new tools for improving crops. *Biol. Res.* **46**, 323–331.
- Fan, D., T. Liu, C. Li, B. Jiao, S. Li, Y. Hou, and K. Luo, 2015: Efficient CRISPR/Cas9-mediated targeted mutagenesis in populus in the first generation. *Sci. Rep.* **5**, 12217.
- Fichtner, F., R. Urrea Castellanos, and B. Ulker, 2014: Precision genetic modifications: a new era in molecular biology and crop improvement. *Planta* **239**, 921–939.
- Gadaleta, A., A. Giancaspro, A. E. Blechl, and A. Blanco, 2008: A transgenic durum wheat line that is free of marker genes and expresses 1Dy10. *J. Cereal Sci.* **48**, 439–445.
- Gaj, T., C. A. Gersbach, and C. F. Barbás 3rd, 2013: ZFN, TALEN, and CRISPR/Cas-based methods for genome engineering. *Trends Biotechnol.* **31**, 397–405.
- Gao, J., G. Wang, S. Ma, X. Xie, X. Wu, X. Zhang, Y. Wu, P. Zhao, and Q. Xia, 2015: CRISPR/Cas9-mediated targeted mutagenesis in *Nicotiana tabacum*. *Plant Mol. Biol.* **87**, 99–110.
- Gruskin, D., 2012: Agbiotech 2.0. *Nat. Biotechnol.* **30**, 211–214.
- Gurushidze, M., G. Hensel, S. Hiekel, S. Schedel, V. Valkov, and J. Kumlehn, 2014: True-breeding targeted gene knock-out in barley using designer TALE-nuclease in haploid cells. *PLoS ONE* **9**, e92046.
- Hamilton, J. P., and C. R. Buell, 2012: Advances in plant genome sequencing. *Plant J.* **70**, 177–190.
- Han, K. M., P. Dharmawardhana, R. S. Arias, C. Ma, V. Busov, and S. H. Strauss, 2011: Gibberellin-associated cisgenes modify growth, stature and wood properties in *Populus*. *Plant Biotechnol. J.* **9**, 162–178.
- Hartung, F., and J. Schiemann, 2014: Precise plant breeding using new genome editing techniques: opportunities, safety and regulation in the EU. *Plant J.* **78**, 742–752.
- Haun, W., A. Coffman, B. M. Clasen, Z. L. Demorest, A. Lowy, E. Ray, A. Retterath, T. Stoddard, A. Juillerat, F. Cedrone, L. Mathis, D. F. Voytas, and F. Zhang, 2014: Improved soybean oil quality by targeted mutagenesis of the fatty acid desaturase 2 gene family. *Plant Biotechnol. J.* **12**, 934–940.
- He, J., X. Zhao, A. Laroche, Z. X. Lu, H. Liu, and Z. Li, 2014: Genotyping-by-sequencing (GBS), an ultimate marker-assisted selection (MAS) tool to accelerate plant breeding. *Front. Plant Sci.* **5**, 484.
- Holme, I. B., G. Dionisio, H. Brinch-Pedersen, T. Wendt, C. K. Madsen, E. Vincze, and P. B. Holm, 2012: Cisgenic barley with improved phytase activity. *Plant Biotechnol. J.* **10**, 237–247.
- Holme, I. B., T. Wendt, and P. B. Holm, 2013: Intragenesis and cisgenesis as alternatives to transgenic crop development. *Plant Biotechnol. J.* **11**, 395–407.
- Hunter, P., 2014: “Genetically Modified Lite” placates public but not activists: new technologies to manipulate plant genomes could help to overcome public concerns about GM crops. *EMBO Rep.* **15**, 138–141.
- Jacobs, T. B., P. R. LaFayette, R. J. Schmitz, and W. A. Parrott, 2015: Targeted genome modifications in soybean with CRISPR/Cas9. *BMC Biotechnol.* **15**, 16.
- Jacobsen, E., and H. J. Schouten, 2007: Cisgenesis strongly improves introgression breeding and induced translocation breeding of plants. *Trends Biotechnol.* **25**, 219–223.
- Jacobsen, E., and H. J. Schouten, 2009: Cisgenesis: an important sub-invention for traditional plant breeding companies. *Euphytica* **170**, 235–247.
- James, C., 2014: Global Status of Commercialized Biotech/GM Crops: 2014 ISAAA Briefs No. 49. ISAAA, Ithaca, NY.
- Jia, H., and N. Wang, 2014: Targeted genome editing of sweet orange using Cas9/sgRNA. *PLoS ONE* **9**, e93806.
- Jiang, W., H. Zhou, H. Bi, M. Fromm, B. Yang, and D. P. Weeks, 2013: Demonstration of CRISPR/Cas9/sgRNA-mediated targeted gene modification in *Arabidopsis*, tobacco, sorghum and rice. *Nucleic Acids Res.* **41**, e188.
- Jinek, M., K. Chylinski, I. Fonfara, M. Hauer, J. A. Doudna, and E. Charpentier, 2012: A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816–821.
- Jo, K. R., C. J. Kim, S. J. Kim, T. Y. Kim, M. Bergervoet, M. A. Jongsma, R. G. Visser, E. Jacobsen, and J. H. Vossen, 2014: Development of late blight resistant potatoes by cisgene stacking. *BMC Biotechnol.* **14**, 50.
- Jones, H. D., 2015: Regulatory uncertainty over genome editing. *Nat. Plants* **1**, 14011.
- Joshi, S. G., J. G. Schaart, R. Groenwold, E. Jacobsen, H. J. Schouten, and F. A. Krens, 2011: Functional analysis and expression profiling of HcrVf1 and HcrVf2 for development of scab resistant cisgenic and intragenic apples. *Plant Mol. Biol.* **75**, 579–591.
- Keurentjes, J. J. B., G. C. Angenent, M. Dicke, V. A. P. M. Dos Santos, J. Molenaar, W. H. van der Putten, P. C. de Rooter, P. C. Struik, and B. P. H. J. Thomma, 2011: Redefining plant systems biology: from cell to ecosystem. *Trends Plant Sci.* **16**, 183–190.
- Keurentjes, J. J. B., J. Molenaar, and B. J. Zwaan, 2013: Predictive modelling of complex agronomic and biological systems. *Plant, Cell Environ.* **36**, 1700–1710.
- Kim, Y. G., and S. Chandrasegaran, 1994: Chimeric restriction endonuclease. *Proc. Natl. Acad. Sci. U S A* **91**, 883–887.
- Kim, Y. G., J. Cha, and S. Chandrasegaran, 1996: Hybrid restriction enzymes: zinc finger fusions to Fok I cleavage domain. *Proc. Natl. Acad. Sci. U S A* **93**, 1156–1160.
- Kumar, V., and M. Jain, 2015: The CRISPR-Cas system for plant genome editing: advances and opportunities. *J. Exp. Bot.* **66**, 47–57.
- Kuzma, J., and A. Kokotovich, 2011: Renegotiating GM crop regulation. *EMBO Rep.* **12**, 883–888.
- Ladics, G. S., A. Bartholomaeus, P. Bregitzer, N. G. Doerrer, A. Gray, T. Holzhauser, M. Jordan, P. Keese, E. Kok, P. Macdonald, W. Parrott, L. Privalle, A. Raybould, S. Y. Rhee, E. Rice, J. Romeis, J. Vaughn, J. M. Wal, and K. Glenn, 2015: Genetic basis and detection of unintended effects in genetically modified crop plants. *Transgenic Res.* **24**, 587–603.
- Lamalakhmi Devi, E., S.K. Chongtham, P. Holeyachi, N. Kousar, M. Singh, C. Behera, R.S. Telem, N.B. Singh, and S.H. Wani, 2013: Cisgenesis and intragenesis: twin sisters for crop improvement. *Res. J. Agric. Forestry Sci.* **1**, 22–26.
- Li, T., B. Liu, M. H. Spalding, D. P. Weeks, and B. Yang, 2012: High-efficiency TALEN-based gene editing produces disease-resistant rice. *Nat. Biotechnol.* **30**, 390–392.
- Li, Z., Z. B. Liu, A. Xing, B. P. Moon, J. P. Koellhoffer, L. Huang, R. T. Ward, E. Clifton, S. C. Falco, and A. M. Cigan, 2015: Cas9-guide RNA directed genome editing in soybean. *Plant Physiol.* **169**, 960–970.
- Liang, Z., K. Zhang, K. Chen, and C. Gao, 2014: Targeted mutagenesis in *Zea mays* using TALENs and the CRISPR/Cas system. *J. Genet. Genomics* **41**, 63–68.
- Lin, T., G. Zhu, J. Zhang, X. Xu, Q. Yu, Z. Zheng, Z. Zhang, Y. Lun, S. Li, X. Wang, Z. Huang, J. Li, C. Zhang, T. Wang, Y. Zhang, A. Wang, Y. Zhang, K. Lin, C. Li, G. Xiong, Y. Xue, A. Mazzucato, M. Causse, Z. Fei, J. J. Giovannoni, R. T. Chetelat, D. Zamir, T. Stadler, J. Li, Z. Ye, Y. Du, and S. Huang, 2014: Genomic analyses provide insights into the history of tomato breeding. *Nat. Genet.* **46**, 1220–1226.
- Liu, L., and X. D. Fan, 2014: CRISPR-Cas system: a powerful tool for genome engineering. *Plant Mol. Biol.* **85**, 209–218.
- Lor, V. S., C. G. Starker, D. F. Voytas, D. Weiss, and N. E. Olszewski, 2014: Targeted mutagenesis of the tomato PROCERA gene using transcription activator-like effector nucleases. *Plant Physiol.* **166**, 1288–1291.
- Lusser, M., and H. V. Davies, 2013: Comparative regulatory approaches for groups of new plant breeding techniques. *New Biotechnol.* **30**, 437–446.
- Lusser, M., C. Parisi, D. Plan, and E. Rodriguez-Cerezo, 2012: Deployment of new biotechnologies in plant breeding. *Nat. Biotechnol.* **30**, 231–239.

- Mahfouz, M. M., A. Piatek, and C. N. Stewart, 2014: Genome engineering via TALENs and CRISPR/Cas9 systems: challenges and perspectives. *Plant Biotechnol. J.* **12**, 1006—1014.
- Mascher, M., T. A. Richmond, D. J. Gerhardt, A. Himmelbach, L. Clissold, D. Sampath, S. Ayling, B. Steuernagel, M. Pfeifer, M. D'Ascenzo, E. D. Akhunov, P. E. Hedley, A. M. Gonzales, P. L. Morrell, B. Kilian, F. R. Blattner, U. Scholz, K. F. Mayer, A. J. Flavell, G. J. Muehlbauer, R. Waugh, J. A. Jeddelloh, and N. Stein, 2013: Barley whole exome capture: a tool for genomic research in the genus *Hordeum* and beyond. *Plant J.* **76**, 494—505.
- Michael, T. P., and R. VanBuren, 2015: Progress, challenges and the future of crop genomes. *Curr. Opin. Plant Biol.* **24**, 71—81.
- Neves, L. G., J. M. Davis, W. B. Barbazuk, and M. Kirst, 2013: Whole-exome targeted sequencing of the uncharacterized pine genome. *Plant J.* **75**, 146—156.
- Nicolia, A., E. Proux-Wera, I. Ahman, N. Onkokesung, M. Andersson, E. Andreasson, and L. H. Zhu, 2015: Targeted gene mutation in tetraploid potato through transient TALEN expression in protoplasts. *J. Biotechnol.* **204**, 17—24.
- Palmgren, M. G., A. K. Edenbrandt, S. E. Vedel, M. M. Andersen, X. Landes, J. T. Osterberg, J. Falhof, L. I. Olsen, S. B. Christensen, P. Sandoe, C. Gamborg, K. Kappel, B. J. Thorsen, and P. Pagh, 2015: Are we ready for back-to-nature crop breeding? *Trends Plant Sci.* **20**, 155—164.
- Peer, R., G. Rivlin, S. Golobovitch, M. Lapidot, A. Gal-On, A. Vainstein, T. Tzfira, and M. A. Flaishman, 2015: Targeted mutagenesis using zinc-finger nucleases in perennial fruit trees. *Planta* **241**, 941—951.
- Podevin, N., Y. Devos, H. V. Davies, and K. M. Nielsen, 2012: Transgenic or not? No simple answer! New biotechnology-based plant breeding techniques and the regulatory landscape. *EMBO Rep.* **13**, 1057—1061.
- Puchta, H., and F. Fauser, 2014: Synthetic nucleases for genome engineering in plants: prospects for a bright future. *Plant J.* **78**, 727—741.
- Ran, F. A., L. Cong, W. X. Yan, D. A. Scott, J. S. Gootenberg, A. J. Kriz, B. Zetsche, O. Shalem, X. Wu, K. S. Makarova, E. V. Koonin, P. A. Sharp, and F. Zhang, 2015: In vivo genome editing using *Staphylococcus aureus* Cas9. *Nature* **520**, 186—191.
- Ricroch, A. E., and M. C. Henard-Damave, 2015: Next biotech plants: new traits, crops, developers and technologies for addressing global challenges. *Crit. Rev. Biotechnol.* **35**, 1—16.
- Rinaldo, A. R., and M. Ayliffe, 2015: Gene targeting and editing in crop plants: a new era of precision opportunities. *Mol. Breeding* **35**, 1—15.
- Rommens, C. M., J. M. Humara, J. Ye, H. Yan, C. Richael, L. Zhang, R. Perry, and K. Swords, 2004: Crop improvement through modification of the plant's own genome. *Plant Physiol.* **135**, 421—431.
- Rommens, C. M., J. S. Ye, C. Richael, and K. Swords, 2006: Improving potato storage and processing characteristics through all-native DNA transformation. *J. Agric. Food Chem.* **54**, 9882—9887.
- Rommens, C. M., H. Yan, K. Swords, C. Richael, and J. S. Ye, 2008: Low-acrylamide French fries and potato chips. *Plant Biotechnol. J.* **6**, 843—853.
- Saintenac, C., D. Jiang, and E. D. Akhunov, 2011: Targeted analysis of nucleotide and copy number variation by exon capture in allotetraploid wheat genome. *Genome Biol.* **12**, R88.
- Sawai, S., K. Ohyama, S. Yasumoto, H. Seki, T. Sakuma, T. Yamamoto, Y. Takebayashi, M. Kojima, H. Sakakibara, T. Aoki, T. Muranaka, K. Saito, and N. Umemoto, 2014: Sterol side chain reductase 2 is a key enzyme in the biosynthesis of cholesterol, the common precursor of toxic steroidal glycoalkaloids in potato. *Plant Cell* **26**, 3763—3774.
- Schaart, J.G., 2004: Towards consumer-friendly cisgenic strawberries which are less susceptible to *Botrytis cinerea*. Ph.D., Wageningen UR, Wageningen, the Netherlands.
- Schaeffer, S. M., and P. A. Nakata, 2015: CRISPR/Cas9-mediated genome editing and gene replacement in plants: transitioning from lab to field. *Plant Sci.* **240**, 130—142.
- Schouten, H., 2014: Reply to Cisgenesis as a golden mean. *Nat. Biotechnol.* **32**, 728.
- Schouten, H. J., F. A. Krens, and E. Jacobsen, 2006: Cisgenic plants are similar to traditionally bred plants: international regulations for genetically modified organisms should be altered to exempt cisgenesis. *EMBO Rep.* **7**, 750—753.
- Schubert, D., and D. Williams, 2006: 'Cisgenic' as a product designation. *Nat. Biotechnol.* **24**, 1327—1329.
- Shan, Q. W., Y. P. Wang, J. Li, Y. Zhang, K. L. Chen, Z. Liang, K. Zhang, J. X. Liu, J. J. Xi, J. L. Qiu, and C. X. Gao, 2013: Targeted genome modification of crop plants using a CRISPR-Cas system. *Nat. Biotechnol.* **31**, 686—688.
- Shan, Q., Y. Zhang, K. Chen, K. Zhang, and C. Gao, 2015: Creation of fragrant rice by targeted knockout of the OsBADH2 gene using TALEN technology. *Plant Biotechnol. J.* **13**, 791—800.
- Shukla, V. K., Y. Doyon, J. C. Miller, R. C. DeKolver, E. A. Moehle, S. E. Worden, J. C. Mitchell, N. L. Arnold, S. Gopalan, X. Meng, V. M. Choi, J. M. Rock, Y. Y. Wu, G. E. Katibah, G. Zhifang, D. McCaskill, M. A. Simpson, B. Blakeslee, S. A. Greenwalt, H. J. Butler, S. J. Hinkley, L. Zhang, E. J. Rebar, P. D. Gregory, and F. D. Urnov, 2009: Precise genome modification in the crop species *Zea mays* using zinc-finger nucleases. *Nature* **459**, 437—441.
- Sprink, T., J. Metje, and F. Hartung, 2015: Plant genome editing by novel tools: TALEN and other sequence specific nucleases. *Curr. Opin. Biotechnol.* **32**, 47—53.
- Sun, X., Z. Hu, R. Chen, Q. Jiang, G. Song, H. Zhang, and Y. Xi, 2015: Targeted mutagenesis in soybean using the CRISPR-Cas9 system. *Sci. Rep.* **5**, 10342.
- Svitashev, S., J. K. Young, C. Schwartz, H. Gao, S. C. Falco, and A. M. Cigan, 2015: Targeted mutagenesis, precise gene editing, and site-specific gene insertion in maize using Cas9 and guide RNA. *Plant Physiol.* **169**, 931—945.
- Townsend, J. A., D. A. Wright, R. J. Winfrey, F. Fu, M. L. Maeder, J. K. Joung, and D. F. Voytas, 2009: High-frequency modification of plant genes using engineered zinc-finger nucleases. *Nature* **459**, 442—445.
- Upadhyay, S.K., J. Kumar, A. Alok, and R. Tuli, 2013: RNA-guided genome editing for target gene mutations in wheat. *G3: Genes - Genomes - Genetics* **3**, 2233—2238.
- Vanblaere, T., I. Szankowski, J. Schaart, H. Schouten, H. Flachowsky, G. A. L. Broggin, and C. Gessler, 2011: The development of a cisgenic apple plant. *J. Biotechnol.* **154**, 304—311.
- Vanblaere, T., H. Flachowsky, C. Gessler, and G. A. L. Broggin, 2014: Molecular characterization of cisgenic lines of apple 'Gala' carrying the Rvi6 scab resistance gene. *Plant Biotechnol. J.* **12**, 2—9.
- Varshney, R. K., R. Terauchi, and S. R. McCouch, 2014: Harvesting the promising fruits of genomics: applying genome sequencing technologies to crop breeding. *PLoS Biol.* **12**, e1001883.
- de Vetten, N., A. M. Wolters, K. Raemakers, I. van der Meer, R. ter Stege, E. Heeres, P. Heeres, and R. Visser, 2003: A transformation method for obtaining marker-free plants of a cross-pollinating and vegetatively propagated crop. *Nat. Biotechnol.* **21**, 439—442.
- Villa, T. C. C., N. Maxted, M. Scholten, and B. Ford-Lloyd, 2005: Defining and identifying crop landraces. *Plant Genet. Res.* **3**, 373—384.
- Voytas, D. F., 2013: Plant genome engineering with sequence-specific nucleases. *Annu. Rev. Plant Biol.* **64**, 327—350.
- Voytas, D. F., and C. Gao, 2014: Precision genome engineering and agriculture: opportunities and regulatory challenges. *PLoS Biol.* **12**, e1001877.
- Wang, Y., X. Cheng, Q. Shan, Y. Zhang, J. Liu, C. Gao, and J. L. Qiu, 2014: Simultaneous editing of three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery mildew. *Nat. Biotechnol.* **32**, 947—951.
- Wang, M., Y. Liu, C. Zhang, J. Liu, X. Liu, L. Wang, W. Wang, H. Chen, C. Wei, X. Ye, X. Li, and J. Tu, 2015a: Gene editing by co-transformation of TALEN and chimeric RNA/DNA oligonucleotides on the rice OsEPSPS gene and the inheritance of mutations. *PLoS ONE* **10**, e0122755.
- Wang, S., S. Zhang, W. Wang, X. Xiong, F. Meng, and X. Cui, 2015b: Efficient targeted mutagenesis in potato by the CRISPR/Cas9 system. *Plant Cell Rep.* **34**, 1473—1476.

- Weeks, J. T., J. Ye, and C. M. Rommens, 2008: Development of an in planta method for transformation of alfalfa (*Medicago sativa*). *Transgenic Res.* **17**, 587—597.
- Wendt, T., P. B. Holm, C. G. Starker, M. Christian, D. F. Voytas, H. Brinch-Pedersen, and I. B. Holme, 2013: TAL effector nucleases induce mutations at a pre-selected location in the genome of primary barley transformants. *Plant Mol. Biol.* **83**, 279—285.
- Wolt, J.D., K. Wang, and B. Yang, 2015: The regulatory status of genome-edited crops. *Plant Biotechnol. J.*, doi: 10.1111/pbi.12444.
- Woo, J. W., J. Kim, S. I. Kwon, C. Corvalan, S. W. Cho, H. Kim, S. G. Kim, S. T. Kim, S. Choe, and J. S. Kim, 2015: DNA-free genome editing in plants with preassembled CRISPR-Cas9 ribonucleoproteins. *Nat. Biotechnol.* **33**, 1162—1164.
- Würdig, J., H. Flachowsky, A. Saß, A. Peil, and M.-V. Hanke, 2015: Improving resistance of different apple cultivars using the Rvi6 scab resistance gene in a cisgenic approach based on the Flp/FRT recombinase system. *Mol. Breeding* **35**, 95.
- Zetsche, B., J.S. Gootenberg, O.O. Abudayyeh, I.M. Slaymaker, K.S. Makarova, P. Essletzbichler, S.E. Volz, J. Joung, J. van der Oost, A. Regev, E.V. Koonin, and F. Zhang, 2015: Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system. *Cell* **163**, 759—771.
- Zeven, A. C., 1998: Landraces: a review of definitions and classifications. *Euphytica* **104**, 127—139.
- Zhang, Y., F. Zhang, X. Li, J. A. Baller, Y. Qi, C. G. Starker, A. J. Bogdanove, and D. F. Voytas, 2013: Transcription activator-like effector nucleases enable efficient plant genome engineering. *Plant Physiol.* **161**, 20—27.
- Zhang, H., J. Zhang, P. Wei, B. Zhang, F. Gou, Z. Feng, Y. Mao, L. Yang, H. Zhang, N. Xu, and J. K. Zhu, 2014: The CRISPR/Cas9 system produces specific and homozygous targeted gene editing in rice in one generation. *Plant Biotechnol. J.* **12**, 797—807.
- Zhou, J., Z. Peng, J. Long, D. Sosso, B. Liu, J.-S. Eom, S. Huang, S. Liu, C. Vera Cruz, W. B. Frommer, F. F. White, and B. Yang, 2015a: Gene targeting by the TAL effector PthXo2 reveals cryptic resistance gene for bacterial blight of rice. *Plant J.* **82**, 632—643.
- Zhou, X., T. B. Jacobs, L.-J. Xue, S. A. Harding, and C.-J. Tsai, 2015b: Exploiting SNPs for biallelic CRISPR mutations in the outcrossing woody perennial *Populus* reveals 4-coumarate: CoA ligase specificity and redundancy. *New Phytol.* **208**, 298—301.